arXiv:0808.2244v4 [hep-th] 22 Oct 2009

NRCPS-HE-56-08

August, 2008

Connection Between
Non-Abelian Tensor Gauge Fields and Open Strings

George Savvidy [*]

Institute of Nuclear Physics,
Demokritos National Research Center
Agia Paraskevi, GR-15310 Athens, Greece

Abstract

We compare the structure of the tree level scattering amplitudes in non-Abelian tensor gauge field theory and in open string theory with Chan-Paton charges. We limit ourselves considering only lower rank tensor fields in both theories. We identify the symmetric and antisymmetric components of the second rank tensor gauge field with the string excitations on the third and forth levels. In the process of this identification we have been selecting only those parts of the tree level scattering amplitudes in the open string theory which have dimensionless coupling constants in four dimensions. It seems that this subclass of tree level scattering amplitudes may provide important information about the structure of the open string theory and most probably it is equivalent to the non-Abelian tensor gauge field theory.

---

[*] savvidy(AT)inp.demokritos.gr

1 *Introduction*

An infinite tower of particles of high spin naturally appears in the spectrum of different string theories. In the low energy limit the massless states of the open string theory with Chan-Paton charges can be identified with the Yang-Mills gauge quanta [1, 2, 3]. It is also expected that in the tensionless limit or, what is equivalent, at high energy and fixed angle scattering the string spectrum becomes effectively massless [4, 5, 6, 7, 15, 16, 8, 9, 17, 10, 11, 12, 13, 18, 14]. It is of great importance to identify these states with the spectrum of some Lagrangian quantum field theory [19, 20, 21, 23, 22, 24, 25].

One can imagine that these massless states are combined into the infinite tower of tensor gauge fields and one could guess that the possible solution could be found by the extension of the Yang-Mills gauge symmetry [26] to non-Abelian tensor gauge fields. This possibility was suggested recently in [27, 28, 29]. Recall that non-Abelian gauge fields are defined as rank-(s+1) tensor gauge fields $A^a_{\mu\lambda_1...\lambda_s}$ [†] and that one can construct infinite series of forms $\mathcal{L}_s$ $(s = 1, 2, ..)$ which are invariant with respect to the extended gauge transformations [27, 28, 29]. These forms $\mathcal{L}_s$ are quadratic in the field strength tensors $G^a_{\mu\nu,\lambda_1...\lambda_s}$ and the general Lagrangian is an infinite sum of these forms (1.1).

The resulting gauge invariant Lagrangian defines *cubic and quartic interactions* with *dimensionless coupling constant* between charged gauge quanta [27, 28, 29]

$$A^a_{\mu}, \qquad A^a_{\mu\lambda_1}, \qquad A^a_{\mu\lambda_1\lambda_2}, \qquad .....$$

carrying a spin larger than one. Note that all these non-Abelian tensor gauge bosons have the same isotopic charges as the vector gauge boson A^a_μ. The gauge invariant Lagrangian describing dynamical tensor gauge bosons of all ranks has the form

$$\mathcal{L} \;=\; \mathcal{L}_{YM} + \mathcal{L}_2 + g_3 \mathcal{L}_3 + ..., \tag{1.1}$$

where $\mathcal{L}_{YM}$ is the Yang-Mills Lagrangian. For the lower-rank tensor gauge fields the Lagrangian has the following form [27, 28, 29]:

$$\mathcal{L}_{YM} = -\frac{1}{4} G^a_{\mu\nu} G^a_{\mu\nu},$$

[†] Tensor gauge fields $A^a_{\mu\lambda_1...\lambda_s}(x)$, $s = 0, 1, 2, ...$ are totally symmetric with respect to the indices $\lambda_1...\lambda_s$. *A priori* the tensor fields have no symmetries with respect to the first index μ. In particular, we have $A^a_{\mu\lambda} \neq A^a_{\lambda\mu}$ and $A^a_{\mu\lambda\rho} = A^a_{\mu\rho\lambda} \neq A^a_{\lambda\mu\rho}$. The adjoint group index $a = 1, ..., N^2 - 1$ in the case of $SU(N)$ gauge group.

$$
\begin{aligned}
\mathcal{L}_2 = &-\frac{1}{4}G^a_{\mu\nu,\lambda}G^a_{\mu\nu,\lambda}-\frac{1}{4}G^a_{\mu\nu}G^a_{\mu\nu,\lambda\lambda} \\
&+\frac{1}{4}G^a_{\mu\nu,\lambda}G^a_{\mu\lambda,\nu}+\frac{1}{4}G^a_{\mu\nu,\nu}G^a_{\mu\lambda,\lambda}+\frac{1}{2}G^a_{\mu\nu}G^a_{\mu\lambda,\nu\lambda},
\end{aligned}
\tag{1.2}
$$

where the generalized field strength tensors are:

$$
\begin{aligned}
G^a_{\mu\nu} &= \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g f^{abc}\, A^b_\mu\, A^c_\nu, \\
G^a_{\mu\nu,\lambda} &= \partial_\mu A^a_{\nu\lambda} - \partial_\nu A^a_{\mu\lambda} + g f^{abc}(\, A^b_\mu\, A^c_{\nu\lambda} + A^b_{\mu\lambda}\, A^c_\nu\,), \\
G^a_{\mu\nu,\lambda\rho} &= \partial_\mu A^a_{\nu\lambda\rho} - \partial_\nu A^a_{\mu\lambda\rho} + g f^{abc}(\, A^b_\mu\, A^c_{\nu\lambda\rho} + A^b_{\mu\lambda}\, A^c_{\nu\rho} + A^b_{\mu\rho}\, A^c_{\nu\lambda} + A^b_{\mu\lambda\rho}\, A^c_\nu\,).
\end{aligned}
\tag{1.3}
$$

The Lagrangian forms $\mathcal{L}_s$ for higher-rank fields can be found in the previous publications [27, 28, 29]. The above expressions define interacting gauge field theory with infinite many non-Abelian tensor gauge fields. Not much is known about physical properties of such gauge field theory and this article is one in the series of articles devoted to its studies [30, 31, 32, 33, 34].

In the present paper we shall focus our attention on the lower-rank tensor gauge field $A^a_{\mu\lambda}$, which decomposes in this theory to the symmetric tensor T_S of helicity two and antisymmetric tensor T_A of helicity zero charged gauge bosons [29]. The Feynman rules for these propagating modes and their interaction vertices can be extracted from the above Lagrangian (1.2) [29] and are reviewed in the next section. These Feynman rules allow, in particular, to calculate tree-level scattering amplitudes for the production of tensor gauge boson in annihilation processes [33, 34].

It is an interesting question if this non-Abelian tensor gauge field theory have any connection with the open string theory with Chan-Paton charges [35][‡]. In the spectrum of the open string theory with Chan-Paton charges there is a massless vector gauge boson V on the second level and there are rank-two massive tensor bosons T_S on the third level and T_A on the forth level carrying the same isotopic charges as the vector boson V. These states are depicted schematically on the Fig.1 as T_S and T_A. The emission vertices for these states are defined as follows [2, 3, 36]:

$$
\begin{aligned}
&e_\alpha(k) : \dot{X}^\alpha e^{ikX} : && \alpha' k^2 = 0 \\
&\varepsilon_{\alpha\alpha'}(k) : \dot{X}^\alpha \dot{X}^{\alpha'} e^{ikX} : && \alpha' k^2 = -1 \\
&\zeta_{\alpha\alpha'}(k)\frac{1}{2} : (\ddot{X}^\alpha \dot{X}^{\alpha'} - \ddot{X}^{\alpha'} \dot{X}^\alpha) e^{ikX} : && \alpha' k^2 = -2.
\end{aligned}
\tag{1.4}
$$

[‡] This question was raised to the author by Costas Bachas and initiated this investigation.



Figure 1: The first excited levels of the open bosonic string. The state at the third level is a symmetric, traceless, rank-2 tensor T_S of $SO(D-1)$. At the fourth excited level there is an antisymmetric rank-2 tensor T_A, of $SO(D-1)$. If the open string carries Chan-Paton charges at its endpoints then these excited states have the same isotopic charges as the massless vector boson V on the first excited level.

They allow to calculate different tree level scattering amplitudes for these tensor bosons.

Our intension in this article is to compare the tree-level scattering amplitudes of the second rank tensor gauge bosons in non-Abelian tensor gauge field theory and tree-level scattering amplitudes of the tensor bosons in open string theory with Chan-Paton charges.

Our aim is two-fold: first, to review the general structure of the interaction vertices in non-Abelian tensor gauge field theory [27, 28, 29] and, second, to calculate similar tree-level scattering amplitudes in the open string theory in order to compare their structures. By "similar" tree-level scattering amplitudes we mean the special selection of amplitudes of the open string theory when the corresponding amplitudes in four-dimensional space-time have *dimensionless coupling constants.* These amplitudes should have only one space-time derivative in the case of triple interaction vertices and have no derivatives in the case of quartic interactions. We are choosing this subclass of interaction amplitudes of open string theory because this subsector of amplitudes with dimensionless coupling constants naturally appears in the low-energy limit when one can ignore the higher derivative terms. Only this subclass of vertices appears in non-Abelian tensor gauge field theory [27, 28, 29].

The open string tree-level scattering amplitudes are defined only on the mass-shell [2, 3]. Therefore in order to compare them with the tensor gauge field theory vertices we have to project them to the mass-shell. We have found that after projection the scattering amplitudes with dimensionless coupling constant for the lower-rank tensor fields in these theories coincide. This result tells that most probably the subsector of the open string theory with dimensionless coupling constants is equivalent to the non-Abelian tensor gauge field theory.

2 *Feynman Rules for Non-Abelian Tensor Gauge Fields*

We shall describe here the interaction vertices of the vector and tensor gauge bosons in non-Abelian tensor gauge field theory [27, 28, 29]. For that let us recapitulate the construction of the corresponding Feynman rules [29]. Because the Lagrangian (1.1) is quadratic in field strength tensors $(G)^2 = (dA + g[A,A])^2$, it allows only *cubic and quartic interactions* with *dimensionless coupling constants*:

$$g\ [A,A]dA, \qquad\qquad g^2\ [A,A][A,A].$$

In particular, the interaction of the Yang-Mills vector bosons with the charged tensor gauge bosons described by the Lagrangian (1.2) are of this type. The Lagrangian (1.2) can be represented in the form of polynomial in Yang-Mills vector field A^a_α and tensor gauge field of the second rank $A^a_{\alpha\acute{\alpha}}$ [29]:

$$\mathcal{L}_2 = \frac{1}{2} A^a_{\alpha\acute{\alpha}} H_{\alpha\acute{\alpha}\gamma\acute{\gamma}} A^a_{\gamma\acute{\gamma}} + \frac{1}{2!}\mathcal{V}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}} A^a_{\alpha\acute{\alpha}} A^b_{\beta} A^c_{\gamma\acute{\gamma}} + \frac{1}{2!2!}\mathcal{V}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}} A^a_{\alpha} A^b_{\beta} A^c_{\gamma\acute{\gamma}} A^d_{\delta\acute{\delta}} + ... \tag{2.5}$$

It has the kinetic term AHA for the tensor gauge field and the interaction vertices between two tensors and a vector, the VTT-vertex $\mathcal{V}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}$ and two tensors and two vectors, the VVTT-vertex $\mathcal{V}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}$. The kinetic operator of the Lagrangian is

$$\begin{aligned} H_{\alpha\acute{\alpha}\gamma\acute{\gamma}}(k) = (-\eta_{\alpha\gamma}\eta_{\acute{\alpha}\acute{\gamma}} + \frac{1}{2}\eta_{\alpha\acute{\gamma}}\eta_{\acute{\alpha}\gamma} + \frac{1}{2}\eta_{\alpha\acute{\alpha}}\eta_{\gamma\acute{\gamma}})k^2 + \eta_{\alpha\gamma}k_{\acute{\alpha}}k_{\acute{\gamma}} + \eta_{\acute{\alpha}\acute{\gamma}}k_{\alpha}k_{\gamma} \\ -\frac{1}{2}(\eta_{\alpha\acute{\gamma}}k_{\acute{\alpha}}k_{\gamma} + \eta_{\acute{\alpha}\gamma}k_{\alpha}k_{\acute{\gamma}} + \eta_{\alpha\acute{\alpha}}k_{\gamma}k_{\acute{\gamma}} + \eta_{\gamma\acute{\gamma}}k_{\alpha}k_{\acute{\alpha}}) \end{aligned} \tag{2.6}$$

and it is symmetric under simultaneous interchange of the indices $\alpha \leftrightarrow \acute{\alpha}$ and $\gamma \leftrightarrow \acute{\gamma}$, but it is not symmetric with respect to a single interchange $\alpha \leftrightarrow \acute{\alpha}$ or $\gamma \leftrightarrow \acute{\gamma}$, because the tensor field $A^a_{\alpha\acute{\alpha}}$ is not a symmetric tensor. It is also fully gauge invariant operator $k_\alpha H_{\alpha\acute{\alpha}\gamma\acute{\gamma}} = 0,\ k_{\acute{\alpha}} H_{\alpha\acute{\alpha}\gamma\acute{\gamma}} = 0$, therefore there is no negative norm states in the spectrum [27, 28, 29]. It describes the propagation of massless particles with helicities two and zero. Indeed, when k_μ is aligned along the third axis, $k_\mu = (k, 0, 0, k)$, the equation

$$H_{\alpha\acute{\alpha}\gamma\acute{\gamma}}(k) f^{\gamma\acute{\gamma}}(k) = 0 \tag{2.7}$$

has three independent solutions of the helicity two and zero

$$\varepsilon^1_{\alpha\acute{\alpha}} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0,0, & 0,0\\ 0,1, & 0,0\\ 0,0, & -1,0\\ 0,0, & 0,0 \end{pmatrix}, \varepsilon^2_{\alpha\acute{\alpha}} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0,0,0,0\\ 0,0,1,0\\ 0,1,0,0\\ 0,0,0,0 \end{pmatrix}, \zeta_{\alpha\acute{\alpha}} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0, & 0,0,0\\ 0, & 0,1,0\\ 0, & -1,0,0\\ 0, & 0,0,0 \end{pmatrix}, \tag{2.8}$$

with the property that $\varepsilon^1_{\gamma\acute{\gamma}}\varepsilon^1_{\lambda\acute{\lambda}} + \varepsilon^2_{\gamma\acute{\gamma}}\varepsilon^2_{\lambda\acute{\lambda}} \simeq \frac{1}{2}(\eta_{\gamma\lambda}\eta_{\acute{\gamma}\acute{\lambda}} + \eta_{\gamma\acute{\lambda}}\eta_{\acute{\gamma}\lambda} - \eta_{\gamma\acute{\gamma}}\eta_{\lambda\acute{\lambda}})$ and $\zeta_{\gamma\acute{\gamma}}\zeta_{\lambda\acute{\lambda}} \simeq \frac{1}{2}(\eta_{\gamma\lambda}\eta_{\acute{\gamma}\acute{\lambda}} - \eta_{\gamma\acute{\lambda}}\eta_{\acute{\gamma}\lambda})$. The symbol $\simeq$ means that the equation holds up to longitudinal terms. The second-rank tensor gauge field $A_{\alpha\acute{\alpha}}$ with 16 components describes in this theory three physical transversal polarizations. The propagator $\Delta_{\gamma\acute{\gamma}\lambda\acute{\lambda}}(k)$ is defined through the equation $H^{fix}_{\alpha\acute{\alpha}\gamma\acute{\gamma}}(k)\Delta^{\gamma\acute{\gamma}}{}_{\lambda\acute{\lambda}}(k) = \eta_{\alpha\lambda}\eta_{\acute{\alpha}\acute{\lambda}}$, and has the following form:

$$\Delta_{\gamma\acute{\gamma}\lambda\acute{\lambda}}(k) = -\frac{4\eta_{\gamma\lambda}\eta_{\gamma'\lambda'} + 2\eta_{\gamma\lambda'}\eta_{\gamma'\lambda} - 3\eta_{\gamma\gamma'}\eta_{\lambda\lambda'}}{3(k^2 - i\varepsilon)} \quad . \tag{2.9}$$

The corresponding residue can be represented as a sum

$$\frac{4\eta_{\gamma\lambda}\eta_{\gamma'\lambda'} + 2\eta_{\gamma\lambda'}\eta_{\gamma'\lambda} - 3\eta_{\gamma\gamma'}\eta_{\lambda\lambda'}}{3} = (\eta_{\gamma\lambda}\eta_{\acute{\gamma}\acute{\lambda}} + \eta_{\gamma\acute{\lambda}}\eta_{\acute{\gamma}\lambda} - \eta_{\gamma\acute{\gamma}}\eta_{\lambda\acute{\lambda}}) + \frac{1}{3}(\eta_{\gamma\lambda}\eta_{\acute{\gamma}\acute{\lambda}} - \eta_{\gamma\acute{\lambda}}\eta_{\acute{\gamma}\lambda}).$$

The first term describes the $\lambda = \pm 2$ helicity states and is represented by the symmetric part $\varepsilon_{\alpha\acute{\alpha}}$ of the polarization tensor, the second term describes $\lambda = 0$ helicity state and is represented by its antisymmetric part $\zeta_{\alpha\acute{\alpha}}$.

Let us now consider three-particle interaction vertex - VTT. Explicitly first three-linear term of the Lagrangian (1.1) has the following form:

$$-\frac{1}{2}gf^{abc}(\partial_\mu A^a_{\nu\lambda} - \partial_\nu A^a_{\mu\lambda})\ (A^b_\mu A^c_{\nu\lambda} + A^b_{\mu\lambda}A^c_\nu) - \frac{1}{4}gf^{abc}(\partial_\mu A^a_\nu - \partial_\nu A^a_\mu)\ 2A^b_{\mu\lambda}A^c_{\nu\lambda}.$$

This is in addition to the standard Yang-Mills VVV three-vector boson interaction vertex

$$\mathcal{L}^{cubic}_1 = -\frac{1}{2}gf^{abc}(\partial_\mu A^a_\nu - \partial_\nu A^a_\mu)A^b_\mu A^c_\nu,$$

which in the momentum representation has the form

$$\mathcal{V}^{abc}_{\alpha\beta\gamma}(k,p,q) = -igf^{abc}F_{\alpha\beta\gamma}(k,p,q) = -igf^{abc}[\eta_{\alpha\beta}(p-k)_\gamma + \eta_{\alpha\gamma}(k-q)_\beta + \eta_{\beta\gamma}(q-p)_\alpha]. \tag{2.10}$$

In momentum space our vertex has the form

$$-igf^{abc}F_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q) = -igf^{abc}[\eta_{\alpha\beta}(p-k)_\gamma + \eta_{\alpha\gamma}(k-q)_\beta + \eta_{\beta\gamma}(q-p)_\alpha]\eta_{\acute{\alpha}\acute{\gamma}}. \tag{2.11}$$



Figure 2: The interaction vertex for the vector gauge boson V and two tensor gauge bosons T - the VTT vertex - $\mathcal{V}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q)$ in non-Abelian tensor gauge field theory [29]. Vector gauge bosons are conventionally drawn as thin wave lines, tensor gauge bosons are thick wave lines. The Lorentz indices $\alpha\acute{\alpha}$ and momentum k belong to the first tensor gauge boson, the $\gamma\acute{\gamma}$ and momentum q belong to the second tensor gauge boson, and Lorentz index β and momentum p belong to the vector gauge boson.

We have also a second part of the three-particle interaction vertex VTT. Explicitly the second three-linear term of Lagrangian (1.1) has the following form:

$$\begin{aligned}
&+\frac{1}{2}gf^{abc}(\partial_\mu A^a_{\nu\lambda}-\partial_\nu A^a_{\mu\lambda})\ (A^b_\mu A^c_{\lambda\nu}+A^b_{\mu\nu}A^c_\lambda)\\
&+\frac{1}{2}gf^{abc}(\partial_\mu A^a_{\nu\nu}-\partial_\nu A^a_{\mu\nu})\ (A^b_\mu A^c_{\lambda\lambda}+A^b_{\mu\lambda}A^c_\lambda)\\
&+\frac{1}{2}gf^{abc}(\partial_\mu A^a_{\nu}-\partial_\nu A^a_{\mu})\ (A^b_{\mu\nu}A^c_{\lambda\lambda}+A^b_{\mu\lambda}A^c_{\lambda\nu}),
\end{aligned}\tag{2.12}$$

so that in the momentum space we have

$$\begin{aligned}
ig\frac{1}{2}f^{abc}F^{'}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q)=ig\frac{1}{2}f^{abc}[&+(p-k)_\gamma(\eta_{\alpha\acute{\gamma}}\eta_{\acute{\alpha}\beta}+\eta_{\alpha\acute{\alpha}}\eta_{\beta\acute{\gamma}})\\
&+(k-q)_\beta(\eta_{\alpha\acute{\gamma}}\eta_{\acute{\alpha}\gamma}+\eta_{\alpha\acute{\alpha}}\eta_{\gamma\acute{\gamma}})\\
&+(q-p)_\alpha(\eta_{\acute{\alpha}\gamma}\eta_{\beta\acute{\gamma}}+\eta_{\acute{\alpha}\beta}\eta_{\gamma\acute{\gamma}})\\
&+(p-k)_{\acute{\alpha}}\eta_{\alpha\beta}\eta_{\gamma\acute{\gamma}}+(p-k)_{\acute{\gamma}}\eta_{\alpha\beta}\eta_{\acute{\alpha}\gamma}\\
&+(k-q)_{\acute{\alpha}}\eta_{\alpha\gamma}\eta_{\beta\acute{\gamma}}+(k-q)_{\acute{\gamma}}\eta_{\alpha\gamma}\eta_{\acute{\alpha}\beta}\\
&+(q-p)_{\acute{\alpha}}\eta_{\beta\gamma}\eta_{\alpha\acute{\gamma}}+(q-p)_{\acute{\gamma}}\eta_{\alpha\acute{\alpha}}\eta_{\beta\gamma}].
\end{aligned}\tag{2.13}$$

Collecting two terms of the three-point vertex VTT together we shall get [29]

$$\mathcal{V}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q)=-igf^{abc}\{F_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}-\frac{1}{2}F^{'}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}\}\equiv-igf^{abc}\mathcal{F}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}\tag{2.14}$$



Figure 3: The quartic vertex with two vector gauge bosons and two tensor gauge bosons - the VVTT vertex - $\mathcal{V}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r)$ in non-Abelian tensor gauge field theory [29]. Vector gauge bosons are conventionally drawn as thin wave lines, tensor gauge bosons are thick wave lines. The Lorentz indices $\gamma\acute{\gamma}$ and momentum q belong to the first tensor gauge boson, $\delta\acute{\delta}$ and momentum r belong to the second tensor gauge boson, the index α and momentum k belong to the first vector gauge boson and Lorentz index β and momentum p belong to the second vector gauge boson.

where the indices $(a, \alpha, \acute{\alpha}, k)$ belongs to the tensor gauge boson, (b, β, p) to the vector gauge boson and $(c, \gamma, \acute{\gamma}, q)$ to the second tensor gauge boson (see Fig.2).

Let us consider now four-particle interaction terms of the Lagrangian (1.2). We have the standard four vector boson interaction vertex VVVV

$$\begin{aligned}\mathcal{V}^{abcd}_{\alpha\beta\gamma\delta}(k,p,q,r) = & -g^2 f^{lac} f^{lbd}(\eta_{\alpha\beta}\eta_{\gamma\delta} - \eta_{\alpha\delta}\eta_{\beta\gamma}) \\ & -g^2 f^{lad} f^{lbc}(\eta_{\alpha\beta}\eta_{\gamma\delta} - \eta_{\alpha\gamma}\eta_{\beta\delta}) \\ & -g^2 f^{lab} f^{lcd}(\eta_{\alpha\gamma}\eta_{\beta\delta} - \eta_{\alpha\delta}\eta_{\beta\gamma}) \end{aligned} \tag{2.15}$$

and a new interaction of two vector and two tensor gauge bosons - the VVTT vertex,

$$\begin{aligned} & -\frac{1}{4}g^2 f^{abc} f^{a\acute{b}\acute{c}}(A^b_\mu A^c_{\nu\lambda} + A^b_{\mu\lambda}A^c_\nu)(A^{\acute{b}}_\mu A^{\acute{c}}_{\nu\lambda} + A^{\acute{b}}_{\mu\lambda}A^{\acute{c}}_\nu) \\ & -\frac{1}{2}g^2 f^{abc} f^{a\acute{b}\acute{c}} A^b_\mu A^c_\nu A^{\acute{b}}_{\mu\lambda} A^{\acute{c}}_{\nu\lambda}, \end{aligned} \tag{2.16}$$

which in the momentum space will take the following form:

$$\begin{aligned} F^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r) = & -g^2 f^{lac} f^{lbd}(\eta_{\alpha\beta}\eta_{\gamma\delta} - \eta_{\alpha\delta}\eta_{\beta\gamma})\eta_{\acute{\gamma}\acute{\delta}} \\ & -g^2 f^{lad} f^{lbc}(\eta_{\alpha\beta}\eta_{\gamma\delta} - \eta_{\alpha\gamma}\eta_{\beta\delta})\eta_{\acute{\gamma}\acute{\delta}} \\ & -g^2 f^{lab} f^{lcd}(\eta_{\alpha\gamma}\eta_{\beta\delta} - \eta_{\alpha\delta}\eta_{\beta\gamma})\eta_{\acute{\gamma}\acute{\delta}}. \end{aligned} \tag{2.17}$$

The second part of the vertex VVTT is:

$$+\frac{1}{4}g^2 f^{abc} f^{a\acute{b}\acute{c}}(A^b_\mu A^c_{\nu\lambda} + A^b_{\mu\lambda}A^c_\nu)(A^{\acute{b}}_\mu A^{\acute{c}}_{\lambda\nu} + A^{\acute{b}}_{\mu\nu}A^{\acute{c}}_\lambda)$$

$$
\begin{aligned}
&+\frac{1}{4}g^2 f^{abc} f^{a\acute{b}\acute{c}}(A^b_\mu A^c_{\nu\nu} + A^b_{\mu\nu}A^c_\nu)(A^{\acute{b}}_\mu A^{\acute{c}}_{\lambda\lambda} + \ A^{\acute{b}}_{\mu\lambda}A^{\acute{c}}_\lambda) \\
&+\frac{1}{2}g^2 f^{abc} f^{a\acute{b}\acute{c}} A^b_\mu A^c_\nu (A^{\acute{b}}_{\mu\nu}A^{\acute{c}}_{\lambda\lambda} + A^{\acute{b}}_{\mu\lambda}A^{\acute{c}}_{\lambda\nu}),
\end{aligned}
\tag{2.18}
$$

which in the momentum representation will take the form

$$
\begin{aligned}
\acute{F}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r) = \frac{1}{2}g^2 f^{lac} f^{lbd} [&+ \eta_{\alpha\beta}(\eta_{\gamma\acute{\delta}}\eta_{\acute{\gamma}\delta} + \eta_{\gamma\acute{\gamma}}\eta_{\delta\acute{\delta}}) \\
&- \eta_{\beta\gamma}(\eta_{\alpha\acute{\delta}}\eta_{\acute{\gamma}\delta} + \eta_{\alpha\acute{\gamma}}\eta_{\delta\acute{\delta}}) \\
&- \eta_{\alpha\delta}(\eta_{\beta\acute{\gamma}}\eta_{\gamma\acute{\delta}} + \eta_{\beta\acute{\delta}}\eta_{\gamma\acute{\gamma}}) \\
&+ \eta_{\gamma\delta}(\eta_{\alpha\acute{\delta}}\eta_{\beta\acute{\gamma}} + \eta_{\alpha\acute{\gamma}}\eta_{\beta\acute{\delta}})] \\
\frac{1}{2}g^2 f^{lad} f^{lbc} [&+ \eta_{\alpha\beta}(\eta_{\gamma\acute{\delta}}\eta_{\acute{\gamma}\delta} + \eta_{\gamma\acute{\gamma}}\eta_{\delta\acute{\delta}}) \\
&- \eta_{\alpha\gamma}(\eta_{\beta\acute{\delta}}\eta_{\acute{\gamma}\delta} + \eta_{\beta\acute{\gamma}}\eta_{\delta\acute{\delta}}) \\
&- \eta_{\beta\delta}(\eta_{\alpha\acute{\gamma}}\eta_{\gamma\acute{\delta}} + \eta_{\alpha\acute{\delta}}\eta_{\gamma\acute{\gamma}}) \\
&+ \eta_{\gamma\delta}(\eta_{\alpha\acute{\gamma}}\eta_{\beta\acute{\delta}} + \eta_{\alpha\acute{\delta}}\eta_{\beta\acute{\gamma}})] \\
\frac{1}{2}g^2 f^{lab} f^{lcd} [&+ \eta_{\alpha\gamma}(\eta_{\beta\acute{\gamma}}\eta_{\delta\acute{\delta}} + \eta_{\beta\acute{\delta}}\eta_{\delta\acute{\gamma}}) \\
&- \eta_{\beta\gamma}(\eta_{\alpha\acute{\gamma}}\eta_{\delta\acute{\delta}} + \eta_{\alpha\acute{\delta}}\eta_{\delta\acute{\gamma}}) \\
&- \eta_{\alpha\delta}(\eta_{\beta\acute{\delta}}\eta_{\gamma\acute{\gamma}} + \eta_{\beta\acute{\gamma}}\eta_{\gamma\acute{\delta}}) \\
&+ \eta_{\beta\delta}(\eta_{\alpha\acute{\delta}}\eta_{\gamma\acute{\gamma}} + \eta_{\alpha\acute{\gamma}}\eta_{\gamma\acute{\delta}})].
\end{aligned}
\tag{2.19}
$$

The total vertex is

$$
\mathcal{V}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r) = F^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r) + \acute{F}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}(k,p,q,r). \tag{2.20}
$$

In summary we have off-mass-shell Yang-Mills vertex VVV (2.10), the new vertex VTT (2.14) together with Yang-Mills vertex VVVV (2.15) and the new vertex VVTT (2.20) (see Fig.2,3).

2.1 *Three-Point Amlitudes*

In order to compare these vertices with the corresponding tree-level amplitudes of the open string theory one should project them to the mass-shell, because the string amplitudes can be computed only on the mass-shell [§]. The three point scattering amplitudes for massless particles are equal to zero for the real momenta (k,p,q), but if one allows complex momenta or a different space-time signature [37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 50, 51, 52], then these matrix elements have nontrivial behavior and we will be able to compare them with the open string tree-level amplitudes.

[§] We shall derive the corresponding string amplitudes in the next section.

Thus multiplying the above VTT vertex (2.14) by the vector wave function $e^{\beta}(p)$ and tensor wave functions $f^{\alpha\acute{\alpha}}(k)$ and $f^{\gamma\acute{\gamma}}(q)$ we shall get the amplitude $\mathcal{V}fef$:

$$\begin{aligned}\mathcal{V}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}_{abc}(k,p,q)|_{mass-shell} = -igf^{abc}[\;& +(k-q)^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha'\gamma'} - \frac{1}{2}\eta^{\alpha\gamma'}\eta^{\alpha'\gamma}) \\ & +(q-p)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha'\gamma'} - \frac{1}{2}\eta^{\beta\gamma'}\eta^{\alpha'\gamma}) \\ & -\frac{1}{2}(q-p)^{\alpha'}(\eta^{\beta\gamma}\eta^{\alpha\gamma'} - \frac{1}{2}\eta^{\beta\gamma'}\eta^{\alpha\gamma}) \\ & +(p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha'\gamma'} - \frac{1}{2}\eta^{\alpha'\beta}\eta^{\alpha\gamma'}) \\ & -\frac{1}{2}(p-k)^{\gamma'}(\eta^{\alpha\beta}\eta^{\alpha'\gamma} - \frac{1}{2}\eta^{\alpha'\beta}\eta^{\alpha\gamma})].\end{aligned} \tag{2.21}$$

Here we have used the transversality of the wave functions (2.8):

$$p_{\beta}e^{\beta}(p) = 0, \quad k_{\alpha}f^{\alpha\acute{\alpha}}(k) = k_{\acute{\alpha}}f^{\alpha\acute{\alpha}}(k) = 0, \quad q_{\gamma}f^{\gamma\acute{\gamma}}(q) = q_{\acute{\gamma}}f^{\gamma\acute{\gamma}}(k) = 0. \tag{2.22}$$

and that they are traceless $trf(k) = trf(q) = 0$. In (2.21) and in the subsequent formulas for amplitudes we shall not show the wave functions. The tensor wave function $f_{\alpha\acute{\alpha}}$ here is a sum of symmetric $\varepsilon_{\alpha\acute{\alpha}}$ and antisymmetric $\zeta_{\alpha\acute{\alpha}}$ parts (2.8).

We shell separate the parts of this vertex which are symmetric - T_S and antisymmetric -T_A with respect to the indices of the tensor field of $f_{\alpha\acute{\alpha}} = \varepsilon_{\alpha\acute{\alpha}} + \zeta_{\alpha\acute{\alpha}}$. The symmetric $\mathcal{V}\varepsilon e\varepsilon$ part of the amplitude is

$$\begin{aligned}\mathcal{V}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}_{abc}(k,p,q)|_{mass-shell} = -i\frac{1}{4}gf^{abc}[\;& +(k-q)^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha'\gamma'} + \eta^{\alpha\gamma'}\eta^{\alpha'\gamma}) \\ & +\frac{1}{4}(q-p)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha'\gamma'} + \eta^{\beta\gamma'}\eta^{\alpha'\gamma}) \\ & +\frac{1}{4}(q-p)^{\alpha'}(\eta^{\beta\gamma}\eta^{\alpha\gamma'} + \eta^{\beta\gamma'}\eta^{\alpha\gamma}) \\ & +\frac{1}{4}(p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha'\gamma'} + \eta^{\alpha'\beta}\eta^{\alpha\gamma'}) \\ & +\frac{1}{4}(p-k)^{\gamma'}(\eta^{\alpha\beta}\eta^{\alpha'\gamma} + \eta^{\alpha'\beta}\eta^{\alpha\gamma})].\end{aligned} \tag{2.23}$$

The antisymmetric $\mathcal{V}\zeta e\zeta$ part of the amplitude is

$$\mathcal{V}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}_{abc}(k,p,q)|_{mass-shell} = -i\frac{3}{4}gf^{abc}[\; +(k-q)^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha'\gamma'} - \eta^{\alpha\gamma'}\eta^{\alpha'\gamma})$$

$$
\begin{aligned}
&+\frac{3}{4}(q-p)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}}-\eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma})\\
&-\frac{3}{4}(q-p)^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}}-\eta^{\beta\gamma^{'}}\eta^{\alpha\gamma})\\
&+\frac{3}{4}(p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}}-\eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}})\\
&-\frac{3}{4}(p-k)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma}-\eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})]. \qquad (2.24)
\end{aligned}
$$

The mixed symmetry part $\mathcal{V}\varepsilon e\zeta$ of the amplitude is

$$
\begin{aligned}
\mathcal{V}_{abc}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q)|_{mass-shell}=-i\frac{3}{16}gf^{abc}[&+(q-p)^{\alpha}(\eta^{\alpha^{'}\gamma^{'}}\eta^{\beta\gamma}-\eta^{\alpha^{'}\gamma}\eta^{\beta\gamma^{'}})\\
&+(q-p)^{\alpha^{'}}(\eta^{\alpha\gamma^{'}}\eta^{\beta\gamma}-\eta^{\alpha\gamma}\eta^{\beta\gamma^{'}})\\
&+(p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}}+\eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\beta})\\
&-(p-k)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma}+\eta^{\alpha\gamma}\eta^{\alpha^{'}\beta})]. \qquad (2.25)
\end{aligned}
$$

The last vertex is symmetric under the interchange $(\alpha \leftrightarrow \alpha^{'})$ and antisymmetric under $(\gamma \leftrightarrow \gamma^{'})$. There is also mixed symmetry part of the vertex which is antisymmetric in $(\alpha \leftrightarrow \alpha^{'})$ and symmetric under $(\gamma \leftrightarrow \gamma^{'})$ the $\mathcal{V}\zeta e\varepsilon$ amplitude:

$$
\begin{aligned}
\mathcal{V}_{abc}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q)|_{mass-shell}=-i\frac{3}{16}gf^{abc}[&+(q-p)^{\alpha}(\eta^{\alpha^{'}\gamma^{'}}\eta^{\beta\gamma}+\eta^{\alpha^{'}\gamma}\eta^{\beta\gamma^{'}})\\
&-(q-p)^{\alpha^{'}}(\eta^{\alpha\gamma^{'}}\eta^{\beta\gamma}+\eta^{\alpha\gamma}\eta^{\beta\gamma^{'}})\\
&+(p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}}-\eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\beta})\\
&+(p-k)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma}-\eta^{\alpha\gamma}\eta^{\alpha^{'}\beta})] \qquad (2.26)
\end{aligned}
$$

One can check that the sum of the above four terms (2.23), (2.24),(2.25) and (2.26) gives the total vertex (2.21).

2.2 *Complex Deformation of Momenta*

The nontrivial three-point amplitudes can be defined if one considers complex momenta or the space-time signature $\eta^{\mu\nu}=(-+-+)$. Then the momenta can be chosen as follows [46]:

$$k_1^{\mu}=(\omega,z,iz,k),\quad k_2^{\mu}=(\omega,-z,-iz,k),\quad k_3^{\mu}=(2\omega,0,0,2k),$$

to fulfill the momentum conservation

$$k_1+k_2=k_3.$$

All massless bosons are on mass-shell $k_1^2 = k_2^2 = k_3^2 = 0, \quad (\omega^2 = k^2)$ and

$$k_1 \cdot k_2 = k_2 \cdot k_3 = k_3 \cdot k_1 = 0.$$

Let us first consider the matrix element VVV for the vector gauge bosons. The polarization vectors are

$$e_1^+ = \frac{1}{\sqrt{2}}(\frac{z}{\omega}, 1, -i, -\frac{z}{k}), \quad e_2^+ = \frac{1}{\sqrt{2}}(-\frac{z}{\omega}, 1, -i, \frac{z}{k}), \quad e_3^- = \frac{1}{\sqrt{2}}(0, 1, i, 0)$$

and are orthogonal to the corresponding momenta

$$k_1 \cdot e_1^+ = 0, \quad k_2 \cdot e_2^+ = 0, \quad k_3 \cdot e_3^- = 0.$$

We can compute now the matrix element using trilinear vertex VVV (2.10):

$$M(+,+,-) = F^{\mu_1\mu_2\mu_3}(k_1, k_2, k_3) e^+_{\mu_1}(k_1) e^+_{\mu_2}(k_2) e^-_{\mu_3}(k_3) =$$
$$= -2e_1^+ \cdot e_2^+ \; k_1 \cdot e_3^- - 2e_2^+ \cdot e_3^- \; k_2 \cdot e_1^+ + 2e_3^- \cdot e_1^+ \; k_3 \cdot e_2^+ = 8\sqrt{2}z. \tag{2.27}$$

And indeed, it is nonzero and grows linearly with momentum deformation z. Using spinor representation of the momenta and polarization vectors [44]

$$k_{a\dot{a}} = \lambda_a \tilde{\lambda}_{\dot{a}}, \quad e^+_{a\dot{a}} = \frac{\mu_a \tilde{\lambda}_{\dot{a}}}{< \mu, \lambda >}, \quad e^-_{a\dot{a}} = \frac{\lambda_a \tilde{\mu}_{\dot{a}}}{[\lambda, \mu]},$$

where

$$\lambda_a = (\sqrt{k^+}, \frac{k_x + ik_y}{\sqrt{k^+}}), \quad \tilde{\lambda}_{\dot{a}} = (\sqrt{k^+}, \frac{k_x - ik_y}{\sqrt{k^+}}), \quad k^+ = k_t + k_z \; ,$$

one can see that

$$< 1, 2 > = < 2, 3 > = < 3, 1 > = 0,$$

but

$$[1, 2] = -4z, \quad [2, 3] = 2\sqrt{2}z, \quad [3, 1] = 2\sqrt{2}z,$$

and

$$M(+,+,-) = -\sqrt{2} \; \frac{[1, 2]^4}{[1, 2][2, 3][3, 1]}.$$

We can calculate now the trilinear vertex VTT matrix element using expressions (2.14), (2.23)

$$M(+2,+1,-2) = \mathcal{F}^{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k_1, k_2, k_3) \varepsilon^+_{\alpha\acute{\alpha}}(k_1) e^+_{\beta}(k_2) \varepsilon^-_{\gamma\acute{\gamma}}(k_3) =$$
$$= 2(k_1 + k_3) \cdot e_2^+ \; \varepsilon_1^+ \cdot \varepsilon_3^- \; + (-k_3 - k_2) \cdot \varepsilon_1^+ \cdot \varepsilon_3^- \cdot e_2^+ \; + (k_2 - k_1) \cdot \varepsilon_3^- \cdot \varepsilon_1^+ \cdot e_2^+$$
$$= 12\sqrt{2}z, \tag{2.28}$$

where $\varepsilon^{+}_{\alpha\acute{\alpha}}(k_1) = e^{+}_{\alpha}(k_1)e^{+}_{\acute{\alpha}}(k_1)$, $\varepsilon^{-}_{\gamma\acute{\gamma}}(k_3) = e^{-}_{\gamma}(k_3)e^{-}_{\acute{\gamma}}(k_3)$. It is nonzero and also grows linearly with momentum deformation z. Using spinor representation we shall get

$$M(+2,+1,-2) = -\frac{3\sqrt{2}}{4}\frac{[1,2]^6}{[1,2][2,3]^3[3,1]}. \tag{2.29}$$

2.3 *The Dual Lagrangian*

For completeness let us also recall expression for the dual Lagrangian which is defined as follows [31, 32]:

$$\begin{aligned} \tilde{\mathcal{L}}_2 = &-\frac{1}{4}\tilde{G}^a_{\mu\nu,\lambda}\tilde{G}^a_{\mu\nu,\lambda} - \frac{1}{4}G^a_{\mu\nu}\tilde{G}^a_{\mu\nu,\lambda\lambda} + \\ &+\frac{1}{4}\tilde{G}^a_{\mu\nu,\lambda}\tilde{G}^a_{\mu\lambda,\nu} + \frac{1}{4}\tilde{G}^a_{\mu\nu,\nu}\tilde{G}^a_{\mu\lambda,\lambda} + \frac{1}{2}G^a_{\mu\nu}\tilde{G}^a_{\mu\lambda,\nu\lambda}, \end{aligned} \tag{2.30}$$

where the dual field strength tensors are:

$$\begin{aligned} \tilde{G}^a_{\mu\nu,\lambda} &= \partial_\mu A^a_{\lambda\nu} - \partial_\nu A^a_{\lambda\mu} + gf^{abc}(\ A^b_\mu\ A^c_{\lambda\nu} + A^b_{\lambda\mu}\ A^c_\nu\ ), \\ \tilde{G}^a_{\mu\nu,\lambda\rho} &= \{\ \partial_\mu(\frac{2}{3}A^a_{\lambda\nu\rho} + \frac{2}{3}A^a_{\rho\nu\lambda} - \frac{1}{3}A^a_{\nu\lambda\rho}) + gf^{abc}\ A^b_\mu\ (\frac{2}{3}A^a_{\lambda\nu\rho} + \frac{2}{3}A^a_{\rho\nu\lambda} - \frac{1}{3}A^a_{\nu\lambda\rho}) + \\ &\quad - \partial_\nu(\frac{2}{3}A^a_{\lambda\mu\rho} + \frac{2}{3}A^a_{\rho\mu\lambda} - \frac{1}{3}A^a_{\mu\lambda\rho}) + gf^{abc}\ (\frac{2}{3}A^a_{\lambda\mu\rho} + \frac{2}{3}A^a_{\rho\mu\lambda} - \frac{1}{3}A^a_{\mu\lambda\rho})\ A^c_\nu\ \} \\ &\quad + gf^{abc}\ (\ A^b_{\lambda\mu}\ A^c_{\rho\nu} + A^b_{\rho\mu}\ A^c_{\lambda\nu}\ ). \end{aligned} \tag{2.31}$$

Here we have a similar polynomial expansion of the dual Lagrangian:

$$\tilde{\mathcal{L}}_2 = \frac{1}{2}A^a_{\alpha\acute{\alpha}}\tilde{H}_{\alpha\acute{\alpha}\gamma\acute{\gamma}}A^a_{\gamma\acute{\gamma}} + \frac{1}{1!2!}\tilde{V}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}A^a_{\alpha\acute{\alpha}}A^b_\beta A^c_{\gamma\acute{\gamma}} + \frac{1}{2!2!}\tilde{V}^{abcd}_{\alpha\beta\gamma\acute{\gamma}\delta\acute{\delta}}A^a_\alpha A^b_\beta A^c_{\gamma\acute{\gamma}}A^d_{\delta\acute{\delta}} + ... \tag{2.32}$$

The kinetic term is identical with the kinetic term (2.6) of the Lagrangian $\mathcal{L}_2$

$$\begin{aligned} \tilde{H}_{\alpha\acute{\alpha}\gamma\acute{\gamma}}(k) = (-\eta_{\alpha\gamma}\eta_{\acute{\alpha}\acute{\gamma}} + \frac{1}{2}\eta_{\alpha\acute{\gamma}}\eta_{\acute{\alpha}\gamma} + \frac{1}{2}\eta_{\alpha\acute{\alpha}}\eta_{\gamma\acute{\gamma}})k^2 + \eta_{\alpha\gamma}k_{\acute{\alpha}}k_{\acute{\gamma}} + \eta_{\acute{\alpha}\acute{\gamma}}k_\alpha k_\gamma \\ -\frac{1}{2}(\eta_{\alpha\acute{\gamma}}k_{\acute{\alpha}}k_\gamma + \eta_{\acute{\alpha}\gamma}k_\alpha k_{\acute{\gamma}} + \eta_{\alpha\acute{\alpha}}k_\gamma k_{\acute{\gamma}} + \eta_{\gamma\acute{\gamma}}k_\alpha k_{\acute{\alpha}}) \end{aligned} \tag{2.33}$$

and is fully gauge invariant operator $k_\alpha\tilde{H}_{\alpha\acute{\alpha}\gamma\acute{\gamma}} = 0$, $k_{\acute{\alpha}}\tilde{H}_{\alpha\acute{\alpha}\gamma\acute{\gamma}} = 0$. One can get also the explicit form of the dual cubic vertex VTT from the dual Lagrangian $\tilde{\mathcal{L}}_2$ (2.30)

$$\tilde{\mathcal{V}}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}} = -igf^{abc}\{\tilde{F}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}} - \frac{1}{2}\tilde{F}'_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}\}, \tag{2.34}$$

where

$$\tilde{F}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q) = [\eta_{\acute{\alpha}\beta}(p-k)_{\acute{\gamma}} + \eta_{\acute{\alpha}\acute{\gamma}}(k-q)_\beta + \eta_{\beta\acute{\gamma}}(q-p)_{\acute{\alpha}}]\eta_{\alpha\gamma} \tag{2.35}$$

and

$$
\begin{aligned}
\tilde{F}'_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}(k,p,q) &= (p-k)_{\acute{\gamma}}(\eta_{\acute{\alpha}\gamma}\eta_{\alpha\beta}+\eta_{\alpha\acute{\alpha}}\eta_{\beta\gamma}) \\
&+ (k-q)_{\beta}(\eta_{\alpha\acute{\gamma}}\eta_{\acute{\alpha}\gamma}+\eta_{\alpha\acute{\alpha}}\eta_{\gamma\acute{\gamma}}) \\
&+ (q-p)_{\acute{\alpha}}(\eta_{\alpha\acute{\gamma}}\eta_{\beta\gamma}+\eta_{\alpha\beta}\eta_{\gamma\acute{\gamma}}) \\
&+ (p-k)_{\alpha}\eta_{\acute{\alpha}\beta}\eta_{\gamma\acute{\gamma}}+(p-k)_{\gamma}\eta_{\acute{\alpha}\beta}\eta_{\alpha\acute{\gamma}} \\
&+ (k-q)_{\alpha}\eta_{\acute{\alpha}\acute{\gamma}}\eta_{\beta\gamma}+(k-q)_{\gamma}\eta_{\acute{\alpha}\acute{\gamma}}\eta_{\alpha\beta} \\
&+ (q-p)_{\alpha}\eta_{\beta\acute{\gamma}}\eta_{\acute{\alpha}\gamma}+(q-p)_{\gamma}\eta_{\alpha\acute{\alpha}}\eta_{\beta\acute{\gamma}}. \qquad (2.36)
\end{aligned}
$$

There is an important property of the dual vertex (2.34) which follows from the fact that the above Lagrangians (1.2) and (2.30) are dual to each other in the sense of the transformation

$$
\begin{aligned}
&\tilde{A}_{\alpha\acute{\alpha}} = A_{\acute{\alpha}\alpha}, \\
&\tilde{A}_{\alpha\acute{\alpha}\acute{\acute{\alpha}}} = \tfrac{2}{3}(A_{\acute{\alpha}\alpha\acute{\acute{\alpha}}}+A_{\acute{\acute{\alpha}}\alpha\acute{\alpha}})-\tfrac{1}{3}A_{\alpha\acute{\acute{\alpha}}}.
\end{aligned} \qquad (2.37)
$$

Indeed, if one simultaneously interchanges the indices $\alpha \leftrightarrow \acute{\alpha}$, $\gamma \leftrightarrow \acute{\gamma}$ of the dual cubic vertex (2.34), then one can see that it will transform into the cubic vertex (2.14) and via versa:

$$
\tilde{\mathcal{V}}^{abc}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}} = -igf^{abc}\{\tilde{F}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}-\frac{1}{2}\tilde{F}'_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}\} = -igf^{abc}\{F_{\acute{\alpha}\alpha\beta\acute{\gamma}\gamma}-\frac{1}{2}F'_{\acute{\alpha}\alpha\beta\acute{\gamma}\gamma}\} = \mathcal{V}^{abc}_{\acute{\alpha}\alpha\beta\acute{\gamma}\gamma} \qquad (2.38)
$$

It is also obvious from the above relation that if one considers the *self-dual sum*

$$
\mathcal{L}_2+\tilde{\mathcal{L}}_2, \qquad (2.39)
$$

then the corresponding VTT vertex will be self-dual in the sense that under duality transformation (2.37) $\alpha \leftrightarrow \acute{\alpha}$, $\gamma \leftrightarrow \acute{\gamma}$ it will be mapped into itself:

$$
\mathcal{V}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}+\tilde{\mathcal{V}}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}} \quad\rightarrow\quad \mathcal{V}_{\acute{\alpha}\alpha\beta\acute{\gamma}\gamma}+\tilde{\mathcal{V}}_{\acute{\alpha}\alpha\beta\acute{\gamma}\gamma} = \tilde{\mathcal{V}}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}+\mathcal{V}_{\alpha\acute{\alpha}\beta\gamma\acute{\gamma}}. \qquad (2.40)
$$

3 *Open Strings Tree-Level Amplitudes*

In this section we shall calculate a tree-level scattering amplitudes in the open string theory with Chan-Paton charges in order to compare them with corresponding matrix elements in non-Abelian tensor gauge field theory. We shall consider scattering amplitudes of the first excited states of open string depicted on the Fig.1, that is the charged scalar, vector and tensor bosons. To set up notation let us begin with the simplest example of the tree-level scattering amplitude for the three on-shell massless vector bosons. The vertex operator has the following form [2, 3]:

$$e_{\alpha}(k) : \dot{X}^{\alpha} e^{ikX}(y) : \tag{3.41}$$

and we shall represent the disk as the upper half-plane so that the boundary coordinate y is real $y \in [-\infty, +\infty]$. The tree amplitude can be represented in the form

$$\mathcal{V}^{\mu_1\mu_2\mu_3}_{a_1a_2a_3}(k_1,k_2,k_3) = F^{\mu_1\mu_2\mu_3}(k_1,k_2,k_3)tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_3}) + F^{\mu_2\mu_1\mu_3}(k_2,k_1,k_3)tr(\lambda^{a_2}\lambda^{a_1}\lambda^{a_3}) :$$

where the matrix element F is given below

$$F^{\mu_1\mu_2\mu_3}(k_1,k_2,k_3) =$$
$$= \int \prod_i d\mu(y_i) <: \dot{X}^{\mu_1} e^{ik_1X}(y_1) :: \dot{X}^{\mu_2} e^{ik_2X}(y_2) :: \dot{X}^{\mu_3} e^{ik_3X}(y_3) :>$$
$$= \lim_{y_1=0, y_2=1, y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \; y_3^2$$
$$\{F^{\mu_1}_{y_1} F^{\mu_2}_{y_2} F^{\mu_2}_{y_2} - 2\alpha' [F^{\mu_1}_{y_1} \frac{\eta^{\mu_2\mu_3}}{(y_2-y_3)^2} + F^{\mu_2}_{y_2} \frac{\eta^{\mu_3\mu_1}}{(y_3-y_1)^2} + F^{\mu_3}_{y_3} \frac{\eta^{\mu_1\mu_2}}{(y_1-y_2)^2}]\}$$

and we have to sum over two orderings of the vertex operators on the disk. The vector functions F^{μ}_{y} are given below (3.44). All bosons are on mass-shell

$$\alpha' k_1^2 = \alpha' k_2^2 = \alpha' k_3^2 = 0$$

and $k_1 + k_2 + k_3 = 0$. The wave functions of the vector bosons are

$$e_{\mu_1}(k_1), \quad e_{\mu_2}(k_2), \quad e_{\mu_3}(k_3) \tag{3.42}$$

and are transversal to the corresponding momenta $k_i \cdot e(k_i) = 0,\ i = 1,2,3$. The matrix element $F^{\mu_1\mu_2\mu_3}$ has the following dimensional structure:

$$[F^{\mu_1\mu_2\mu_3}] \sim (\alpha')^3(k)^3 + (\alpha')^2(k)^1. \tag{3.43}$$

Our intension is to extract that part of the amplitude which has dimensionless coupling constant, that is, the last term. We shall fix the integration measure by the choice $y_1 = 0,\ y_2 = 1,\ y_3 \to \infty$:

$$\lim_{y_1=0,\ y_2=1,\ y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \;\to 1$$

$$
\begin{aligned}
F^{\mu_1}_{y_1} &= -2i\alpha^{'}\ (\frac{k_2^{\mu_1}}{y_1 - y_2} + \frac{k_3^{\mu_1}}{y_1 - y_3}) \rightarrow -2i\alpha^{'}\ (-k_2^{\mu_1}) \\
F^{\mu_2}_{y_2} &= -2i\alpha^{'}\ (\frac{k_1^{\mu_2}}{y_2 - y_1} + \frac{k_3^{\mu_2}}{y_2 - y_3}) \rightarrow -2i\alpha^{'}\ (+k_1^{\mu_2}) \\
F^{\mu_3}_{y_3} &= -2i\alpha^{'}\ (\frac{k_1^{\mu_3}}{y_3 - y_1} + \frac{k_2^{\mu_3}}{y_3 - y_2}) \rightarrow -2i\alpha^{'}\ (-\frac{k_1^{\mu_3}}{y_3^2})
\end{aligned}
\tag{3.44}
$$

Thus for the $F^{\mu_1\mu_2\mu_3}(k_1, k_2, k_3) tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_3})$ we have

$$
i(2\alpha^{'})^2[-k_2^{\mu_1}\eta^{\mu_2\mu_3} + k_1^{\mu_2}\eta^{\mu_3\mu_1} - k_1^{\mu_3}\eta^{\mu_1\mu_2} + 2\alpha^{'}k_2^{\mu_1}k_1^{\mu_2}k_1^{\mu_3}]\ \ tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_3}) \tag{3.45}
$$

and adding the equal term

$$
i(2\alpha^{'})^2[+k_3^{\mu_1}\eta^{\mu_2\mu_3} - k_3^{\mu_2}\eta^{\mu_3\mu_1} + k_2^{\mu_3}\eta^{\mu_1\mu_2} + 2\alpha^{'}k_2^{\mu_1}k_1^{\mu_2}k_1^{\mu_3}]\ \ tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_3}) \tag{3.46}
$$

we can get the total matrix element together with the reversed cyclic orientation $a_1, \mu_1, k_1 \leftrightarrow a_2, \mu_2, k_2$:

$$
\begin{aligned}
2i(\alpha^{'})^2[\ (k_3 - k_2)^{\mu_1}\eta^{\mu_2\mu_3} + (k_1 - k_3)^{\mu_2}\eta^{\mu_3\mu_1} + (k_2 - k_1)^{\mu_3}\eta^{\mu_1\mu_2} + \\
+ 2\alpha^{'}(k_2 - k_3)^{\mu_1}k_1^{\mu_2}k_1^{\mu_3}\ ]\ \ tr([\lambda^{a_1}, \lambda^{a_2}]\lambda^{a_3}).
\end{aligned}
\tag{3.47}
$$

Leaving only dimensionless coupling constant we shall have the corresponding VVV amplitude:

$$
\begin{aligned}
&\mathcal{V}^{\mu_1\mu_2\mu_3}_{a_1a_2a_3}(k_1, k_2, k_3) = \\
&= ig\ tr([\lambda^{a_1}, \lambda^{a_2}]\lambda^{a_3})[\ (k_3 - k_2)^{\mu_1}\eta^{\mu_2\mu_3} + (k_1 - k_3)^{\mu_2}\eta^{\mu_3\mu_1} + (k_2 - k_1)^{\mu_3}\eta^{\mu_1\mu_2}\ ]\ .
\end{aligned}
\tag{3.48}
$$

which coincides with the Yang-Mills vertex (2.10) projected to the mass-shell. In the next subsection we shall perform a similar calculation of the scattering amplitude between two symmetric tensor bosons T_S and a vector boson V in open string theory (see Fig.1) in order to compare it with the amplitude (2.23), (2.28) in non-Abelian tensor gauge field theory.

3.1 *Tree-Level Amplitude for Two Symmetric Tensors and Vector*

The vertex operator for the symmetric T_S rank-2 tensor boson on the third level is

$$
\varepsilon_{\alpha\alpha'}(k) : \dot{X}^{\alpha}\dot{X}^{\alpha^{'}} e^{ikX}(y) : \tag{3.49}
$$

and together with the vertex (3.41) can be used to calculate now the scattering amplitude between vector and two tensor bosons:

$$
\mathcal{V}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{abc}(k, p, q) = F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k, p, q)\ tr(\lambda^a\lambda^b\lambda^c) + F^{\gamma\gamma^{'}\beta\alpha\alpha^{'}}(q, p, k)\ tr(\lambda^c\lambda^b\lambda^a) \tag{3.50}
$$

where the wave functions of the pair of tensor gauge bosons and the vector boson are:

$$\varepsilon_{\alpha\alpha'}(k), \quad e_{\beta}(p), \quad \varepsilon_{\gamma\gamma'}(q). \tag{3.51}$$

We shall define for convenience $k_1 \equiv k, k_2 \equiv q, k_3 \equiv p$, and $k_1+k_2+k_3=0$. The mass-shell conditions are

$$\alpha' k_1^2 = \alpha' k_2^2 = -1, \;\; \alpha' k_3^2 = 0.$$

We have to calculate the correlation function:

$$\begin{aligned}
&F_S^{\alpha\alpha'\beta\gamma\gamma'}(k,p,q) = \\
&= \int \prod_i d\mu(y_i) <: \dot{X}^{\alpha}\dot{X}^{\alpha'} e^{ikX}(y_1) :: \dot{X}^{\gamma}\dot{X}^{\gamma'} e^{iqX}(y_2) :: \dot{X}^{\beta} e^{ipX}(y_3) :> \\
&= \lim_{y_1=0, y_2=1, y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \; y_3^2 \\
&\{ \, (F^{\alpha}F^{\alpha'})_{y_1} F^{\beta}_{y_3} (F^{\gamma}F^{\gamma'})_{y_2} - \\
&- 2\alpha' [F^{\alpha}_{y_1}F^{\beta}_{y_3}F^{\gamma}_{y_2}\frac{\eta^{\alpha'\gamma'}}{(y_1-y_2)^2} + F^{\alpha}_{y_1}F^{\beta}_{y_3}F^{\gamma'}_{y_2}\frac{\eta^{\alpha'\gamma}}{(y_1-y_2)^2} + F^{\alpha'}_{y_1}F^{\beta}_{y_3}F^{\gamma}_{y_2}\frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2} + F^{\alpha'}_{y_1}F^{\beta}_{y_3}F^{\gamma'}_{y_2}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2} + \\
&+ F^{\alpha}_{y_1}F^{\alpha'}_{y_1}F^{\gamma}_{y_2}\frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2} + F^{\alpha}_{y_1}F^{\alpha'}_{y_1}F^{\gamma'}_{y_2}\frac{\eta^{\beta\gamma}}{(y_3-y_2)^2} + F^{\alpha}_{y_1}F^{\gamma}_{y_2}F^{\gamma'}_{y_2}\frac{\eta^{\beta\alpha'}}{(y_3-y_1)^2} + F^{\alpha'}_{y_1}F^{\gamma}_{y_2}F^{\gamma'}_{y_2}\frac{\eta^{\beta\alpha}}{(y_3-y_1)^2}] + \\
&+ (2\alpha')^2 [+F^{\beta}_{y_3}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}\frac{\eta^{\alpha'\gamma'}}{(y_1-y_2)^2} + F^{\beta}_{y_3}\frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2}\frac{\eta^{\alpha'\gamma}}{(y_1-y_2)^2} + \\
&+ F^{\alpha}_{y_1}\frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{\eta^{\alpha'\gamma'}}{(y_1-y_2)^2} + F^{\alpha}_{y_1}\frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{\eta^{\alpha'\gamma}}{(y_1-y_2)^2} + \\
&+ F^{\alpha'}_{y_1}\frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2} + F^{\alpha'}_{y_1}\frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2} + \\
&+ F^{\gamma}_{y_2}\frac{\eta^{\beta\alpha}}{(y_3-y_1)^2}\frac{\eta^{\alpha'\gamma'}}{(y_1-y_2)^2} + F^{\gamma}_{y_2}\frac{\eta^{\beta\alpha'}}{(y_3-y_1)^2}\frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2} + \\
&+ F^{\gamma'}_{y_2}\frac{\eta^{\beta\alpha}}{(y_3-y_1)^2}\frac{\eta^{\alpha'\gamma}}{(y_1-y_2)^2} + F^{\gamma'}_{y_2}\frac{\eta^{\beta\alpha'}}{(y_3-y_1)^2}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}]\}.
\end{aligned} \tag{3.52}$$

The amplitude has the following dimensional structure:

$$F^{\alpha\alpha'\beta\gamma\gamma'} \sim (\alpha')^5(k)^5 + (\alpha')^4(k)^3 + (\alpha')^3(k)^1 \;, \tag{3.53}$$

where the first term contains the fifth power of α' and the fifth power of momentum and so on. *We are interested only in the last term which contains the first power of momentum* *and therefore has the dimensionless coupling constant.* We shall fix the integration measure by the choice $y_1 = 0,\ y_2 = 1,\ y_3 \to \infty$:

$$\lim_{y_1=0,\ y_2=1,\ y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha^{'} k_i k_j} \ \to 1,$$

$$F^{\alpha}_{y_1} = -2i\alpha^{'}\ (\frac{p^{\alpha}}{y_1 - y_3} + \frac{q^{\alpha}}{y_1 - y_2}) \to -2i\alpha^{'}\ (-q^{\alpha})$$

$$F^{\gamma}_{y_2} = -2i\alpha^{'}\ (\frac{p^{\gamma}}{y_2 - y_3} + \frac{k^{\gamma}}{y_2 - y_1}) \to -2i\alpha^{'}\ (k^{\gamma})$$

$$F^{\beta}_{y_3} = -2i\alpha^{'}\ (\frac{k^{\beta}}{y_3 - y_1} + \frac{q^{\beta}}{y_3 - y_2}) \to -2i\alpha^{'}\ (-\frac{k^{\beta}}{y_3^2})\ , \tag{3.54}$$

and keeping only part of the amplitude linear in momentum we shall get:

$$\begin{aligned} F_S^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -i(2\alpha^{'})^3[& -k^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & - q^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & - q^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\ & + k^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\ & + k^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})]. \end{aligned} \tag{3.55}$$

We can add an equal term

$$\begin{aligned} F_S^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -i(2\alpha^{'})^3[& + q^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & + p^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & + p^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\ & - p^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\ & - p^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})] \end{aligned} \tag{3.56}$$

in order to get symmetric expression:

$$\begin{aligned} F_S^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -\frac{i}{2}(2\alpha^{'})^3[& + (q-k)^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & + (p-q)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & + (p-q)^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\ & + (k-p)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\ & + (k-p)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} + \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})]. \end{aligned} \tag{3.57}$$

Substituting this into the expression (3.50) with the terms in the reversed cyclic orientation $a, (\alpha, \alpha^{'}), k \leftrightarrow c, (\gamma, \gamma^{'}), q$ we shall get:

$$\mathcal{V}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{abc}(k,p,q) = \ tr([\lambda^a, \lambda^b]\lambda^c) F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{S}(k,p,q)\ . \tag{3.58}$$

This expression should be compared with the expression (2.23) in tensor gauge field theory. We see that they have a similar Lorentz structure, but with some differences in the coefficients. We don't know exactly the origin of this difference, but most probably it is connected with contributions of higher rank non-Abelian tensor gauge fields, which we do not take into consideration in this article.

3.2 *Tree-Level Amplitude for Two Anti-Symmetric Tensors and Vector*

The vertex operator for the antisymmetric T_A rank-2 tensor boson on the forth level is:

$$\zeta_{\alpha\alpha'}(k) : \ddot{X}^{\{\alpha}\dot{X}^{\alpha^{'}\}} e^{ikX} = \zeta_{\alpha\alpha'}(k)\frac{1}{2} : (\ddot{X}^{\alpha}\dot{X}^{\alpha^{'}} - \ddot{X}^{\alpha^{'}}\dot{X}^{\alpha}) e^{ikX} : . \tag{3.59}$$

The antisymmetric wave functions of the pair of tensor gauge bosons and the vector boson are:

$$\zeta_{\alpha\alpha'}(k), \quad \zeta_{\gamma\gamma'}(q), \quad e_{\beta}(p) \tag{3.60}$$

and we shall define for convenience $k_1 \equiv k, k_2 \equiv q, k_3 \equiv p,$. The mass-shell conditions are

$$\alpha^{'} k_1^2 = \alpha^{'} k_2^2 = -2, \ \alpha^{'} k_3^2 = 0$$

and $k_1 + k_2 + k_3 = 0$. We have to calculate the correlation function:

$$F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{A}(k,p,q) = \tag{3.61}$$

$$= \int \prod_i \ d\mu(y_i) <: \ddot{X}^{\{\alpha}\dot{X}^{\alpha^{'}\}} e^{ikX}(y_1) :: \ddot{X}^{\{\gamma}\dot{X}^{\gamma^{'}\}} e^{iqX}(y_2) :: \dot{X}^{\beta} e^{ipX}(y_3) :> \tag{3.62}$$

$$= \lim_{y_1=0, y_2=1, y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha^{'} k_i k_j} \ y_3^2 \ \frac{1}{4} \quad \{\mathcal{O}(\alpha^{'})^5 (k)^5 + \mathcal{O}(\alpha^{'})^4 (k)^3 +$$

$$+ (2\alpha^{'})^2 F^{\beta}_{y_3} [+ \frac{-6\eta^{\alpha\gamma}}{(y_1-y_2)^4} \frac{\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^2} + \frac{-2\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^3} \frac{2\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^3}$$

$$- \frac{-2\eta^{\alpha\gamma}}{(y_1-y_2)^3} \frac{2\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^3} - \frac{-6\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^4} \frac{\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^2}$$

$$
\begin{aligned}
&\qquad - \frac{2\eta^{\alpha\gamma}}{(y_1-y_2)^3}\frac{-2\eta^{\alpha'\gamma'}}{(y_1-y_2)^3} - \frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2}\frac{-6\eta^{\alpha'\gamma}}{(y_1-y_2)^4} \\
&\qquad + \frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}\frac{-6\eta^{\alpha'\gamma'}}{(y_1-y_2)^4} + \frac{2\eta^{\alpha\gamma'}}{(y_1-y_2)^3}\frac{-2\eta^{\alpha'\gamma}}{(y_1-y_2)^3}] \\
&+ F^{\alpha}_{y_1}[- \frac{2\eta^{\beta\gamma}}{(y_3-y_2)^3}\frac{-2\eta^{\alpha'\gamma'}}{(y_1-y_2)^3} - \frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{-6\eta^{\alpha'\gamma}}{(y_1-y_2)^4} \\
&\qquad + \frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{-6\eta^{\alpha'\gamma'}}{(y_1-y_2)^4} + \frac{2\eta^{\beta\gamma'}}{(y_3-y_2)^3}\frac{-2\eta^{\alpha'\gamma}}{(y_1-y_2)^3}] \\
&+ \dot{F}^{\alpha}_{y_1}[+ \frac{2\eta^{\beta\gamma}}{(y_3-y_2)^3}\frac{\eta^{\alpha'\gamma'}}{(y_1-y_2)^2} + \frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{2\eta^{\alpha'\gamma}}{(y_1-y_2)^3} \\
&\qquad - \frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{2\eta^{\alpha'\gamma'}}{(y_1-y_2)^3} + \frac{2\eta^{\beta\gamma'}}{(y_3-y_2)^3}\frac{\eta^{\alpha'\gamma}}{(y_1-y_2)^2}] \\
&+ F^{\alpha'}_{y_1}[+ \frac{2\eta^{\beta\gamma}}{(y_3-y_2)^3}\frac{-2\eta^{\alpha\gamma'}}{(y_1-y_2)^3} + \frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{-6\eta^{\alpha\gamma}}{(y_1-y_2)^4} \\
&\qquad - \frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{-6\eta^{\alpha\gamma'}}{(y_1-y_2)^4} - \frac{2\eta^{\beta\gamma'}}{(y_3-y_2)^3}\frac{-2\eta^{\alpha\gamma}}{(y_1-y_2)^3}] \\
&+ \dot{F}^{\alpha'}_{y_1}[- \frac{2\eta^{\beta\gamma}}{(y_3-y_2)^3}\frac{\eta^{\alpha\gamma'}}{(y_1-y_2)^2} - \frac{\eta^{\beta\gamma'}}{(y_3-y_2)^2}\frac{2\eta^{\alpha\gamma}}{(y_1-y_2)^3} \\
&\qquad + \frac{\eta^{\beta\gamma}}{(y_3-y_2)^2}\frac{2\eta^{\alpha\gamma'}}{(y_1-y_2)^3} + \frac{2\eta^{\beta\gamma'}}{(y_3-y_2)^3}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}] \\
&+ F^{\gamma}_{y_2}[\ \dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\ ] \\
&+ \dot{F}^{\gamma}_{y_2}[\ \dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\ ] \\
&+ F^{\gamma'}_{y_2}[\ \dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\ ] \\
&+ \dot{F}^{\gamma'}_{y_2}[\ \dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\dots\ ] \quad \},
\end{aligned}
\tag{3.63}
$$

where dots denote the terms which one can get by interchanging α, α' with γ, γ'. This amplitude has the same dimensional structure as the symmetric one:

$$
F^{\alpha\alpha'\beta\gamma\gamma'} \sim (\alpha')^5(k)^5 + (\alpha')^4(k)^3 + (\alpha')^3(k)^1 \tag{3.64}
$$

and we shall calculate the term which contains only the first power of momentum, that is, the last term. Taking the corresponding limit

$$
\lim_{y_1=0,\ y_2=1,\ y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \to 1
$$

$$
F^{\alpha}_{y_1} = -2i\alpha' \left(\frac{p^{\alpha}}{y_1-y_3} + \frac{q^{\alpha}}{y_1-y_2}\right) \to -2i\alpha' (-q^{\alpha})
$$

$$\begin{aligned}
F^{\gamma}_{y_2} &= -2i\alpha^{'}\ (\frac{p^{\gamma}}{y_2-y_3}+\frac{k^{\gamma}}{y_2-y_1}) \rightarrow -2i\alpha^{'}\ (k^{\gamma}) \\
F^{\beta}_{y_3} &= -2i\alpha^{'}\ (\frac{k^{\beta}}{y_3-y_1}+\frac{q^{\beta}}{y_3-y_2}) \rightarrow -2i\alpha^{'}\ (-\frac{k^{\beta}}{y_3^2}) \\
\dot{F}^{\alpha}_{y_1} &= 2i\alpha^{'}\ (\frac{p^{\alpha}}{(y_1-y_3)^2}+\frac{q^{\alpha}}{(y_1-y_2)^2}) \rightarrow -2i\alpha^{'}\ (-q^{\alpha}) \\
\dot{F}^{\gamma}_{y_2} &= 2i\alpha^{'}\ (\frac{p^{\gamma}}{(y_2-y_3)^2}+\frac{k^{\gamma}}{(y_2-y_1)^2}) \rightarrow -2i\alpha^{'}\ (-k^{\gamma})
\end{aligned} \tag{3.65}$$

we shall get

$$\begin{aligned}
F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{A}(k,p,q) = -i(2\alpha^{'})^3[&+ k^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&+ q^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&- q^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\
&+ k^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\
&- k^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})].
\end{aligned} \tag{3.66}$$

Adding the equal term

$$\begin{aligned}
F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{A}(k,p,q) = -i(2\alpha^{'})^3[&- q^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&- p^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&+ p^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\
&- p^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\
&+ p^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})]
\end{aligned} \tag{3.67}$$

we shall get a symmetric in momenta expression

$$\begin{aligned}
F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{A}(k,p,q) = -\frac{i}{2}(2\alpha^{'})^3[&- (q-k)^{\beta}(\eta^{\alpha\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&- (p-q)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\
&+ (p-q)^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha\gamma}) \\
&- (p-k)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma^{'}}) \\
&+ (p-k)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} - \eta^{\alpha^{'}\beta}\eta^{\alpha\gamma})].
\end{aligned} \tag{3.68}$$

Substituting this into the expression (3.50) with the terms in the reversed cyclic orientation $a,(\alpha,\alpha^{'}),k \leftrightarrow c,(\gamma,\gamma^{'}),q$ we shall finally get:

$$\mathcal{V}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{abc}(k,p,q) = tr([\lambda^a,\lambda^b]\lambda^c)F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{A}(k,p,q). \tag{3.69}$$

This expression should be compared with the expression (2.24) and again we see that they have a similar structure.

3.3 *Mixed Symmetry Amplitudes*

Finally we shall calculate the mixed amplitude between symmetric and antisymmetric tensor bosons and a vector. The vertex operator for the symmetric T_S and antisymmetric T_A rank-2 tensor bosons have been given above (3.49) and (3.59). Therefore the matrix element is:

$$\begin{aligned}
&F^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}_{SA}(k,p,q) = \\
&= \int \prod_i \, d\mu(y_i) <: \dot{X}^{\alpha}\dot{X}^{\alpha^{'}} e^{ikX}(y_1) :: \ddot{X}^{\{\gamma}\dot{X}^{\gamma^{'}\}} e^{iqX}(y_2) :: \dot{X}^{\beta} e^{ipX}(y_3) :> \\
&= \lim_{y_1=0, y_2=1, y_3\to\infty} \prod_{i<j} |y_i - y_j|^{2\alpha^{'}k_i k_j} \; y_3^2 \; \frac{1}{2} \quad \{\mathcal{O}(\alpha^{'})^5(k)^5 + \mathcal{O}(\alpha^{'})^4(k)^3 + \\
&+ (2\alpha^{'})^2 F^{\beta}_{y_3}[+\frac{2\eta^{\alpha\gamma}}{(y_1-y_2)^3}\frac{\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^2} + \frac{\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^2}\frac{2\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^3} \\
&\qquad - \frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}\frac{2\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^3} - \frac{2\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^3}\frac{\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^2}] \\
&\quad + F^{\alpha}_{y_1}[+\frac{-2\eta^{\beta\gamma}}{(y_2-y_3)^3}\frac{\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^2} + \frac{\eta^{\beta\gamma^{'}}}{(y_2-y_3)^2}\frac{2\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^3} \\
&\qquad - \frac{\eta^{\beta\gamma}}{(y_2-y_3)^2}\frac{2\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^3} - \frac{-2\eta^{\beta\gamma^{'}}}{(y_2-y_3)^3}\frac{\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^2}] \\
&\quad + F^{\alpha^{'}}_{y_1}[+\frac{-2\eta^{\beta\gamma}}{(y_2-y_3)^3}\frac{\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^2} + \frac{\eta^{\beta\gamma^{'}}}{(y_2-y_3)^2}\frac{2\eta^{\alpha\gamma}}{(y_1-y_2)^3} \\
&\qquad - \frac{\eta^{\beta\gamma}}{(y_2-y_3)^2}\frac{2\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^3} - \frac{-2\eta^{\beta\gamma^{'}}}{(y_2-y_3)^3}\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}] \\
&\quad + \dot{F}^{\gamma}_{y_2}[+\frac{\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^2}\frac{\eta^{\alpha^{'}\beta}}{(y_1-y_3)^2} + \frac{\eta^{\alpha\beta}}{(y_1-y_3)^2}\frac{\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^2}] \\
&\quad + F^{\gamma}_{y_2}[-\frac{2\eta^{\alpha\gamma^{'}}}{(y_1-y_2)^3}\frac{\eta^{\alpha^{'}\beta}}{(y_1-y_3)^2} - \frac{\eta^{\alpha\beta}}{(y_1-y_3)^2}\frac{2\eta^{\alpha^{'}\gamma^{'}}}{(y_1-y_2)^3}] \\
&\quad + F^{\gamma^{'}}_{y_2}[+\frac{2\eta^{\alpha\gamma}}{(y_1-y_2)^3}\frac{\eta^{\alpha^{'}\beta}}{(y_1-y_3)^2} + \frac{\eta^{\alpha\beta}}{(y_1-y_3)^2}\frac{2\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^3}] \\
&\qquad + \dot{F}^{\gamma^{'}}_{y_2}[-\frac{\eta^{\alpha\gamma}}{(y_1-y_2)^2}\frac{\eta^{\alpha^{'}\beta}}{(y_1-y_3)^2} - \frac{\eta^{\alpha\beta}}{(y_1-y_3)^2}\frac{\eta^{\alpha^{'}\gamma}}{(y_1-y_2)^2}]\ \}.
\end{aligned} \tag{3.70}$$

Taking the corresponding limit we shall get

$$\begin{aligned} F_{S\;\;\;\;\;A}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -i(2\alpha^{'})^{3}[\; & - q^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & - q^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} - \eta^{\alpha\gamma}\eta^{\beta\gamma^{'}}) \\ & + \frac{1}{2}k^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\beta}) \\ & - \frac{1}{2}k^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} + \eta^{\alpha\gamma}\eta^{\alpha^{'}\beta})] \end{aligned} \tag{3.71}$$

or, in equivalent form, as

$$\begin{aligned} F_{S\;\;\;\;\;A}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -\frac{i}{2}(2\alpha^{'})^{3}[\; & + (p-q)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & + (p-q)^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} - \eta^{\alpha\gamma}\eta^{\beta\gamma^{'}}) \\ & + \frac{1}{2}(k-p)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\beta}) \\ & - \frac{1}{2}(k-p)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} + \eta^{\alpha\gamma}\eta^{\alpha^{'}\beta})]. \end{aligned} \tag{3.72}$$

The mixed symmetry vertex will take the following form:

$$\begin{aligned} F_{A\;\;\;\;\;S}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = -\frac{i}{2}(2\alpha^{'})^{3}[\; & + \frac{1}{2}(p-q)^{\alpha}(\eta^{\beta\gamma}\eta^{\alpha^{'}\gamma^{'}} + \eta^{\beta\gamma^{'}}\eta^{\alpha^{'}\gamma}) \\ & - \frac{1}{2}(p-q)^{\alpha^{'}}(\eta^{\beta\gamma}\eta^{\alpha\gamma^{'}} + \eta^{\alpha\gamma}\eta^{\beta\gamma^{'}}) \\ & + (k-p)^{\gamma}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma^{'}} - \eta^{\alpha\gamma^{'}}\eta^{\alpha^{'}\beta}) \\ & + (k-p)^{\gamma^{'}}(\eta^{\alpha\beta}\eta^{\alpha^{'}\gamma} - \eta^{\alpha\gamma}\eta^{\alpha^{'}\beta})] \end{aligned} \tag{3.73}$$

Plugging both expressions into (3.50) with the terms in the reversed cyclic orientation $a,(\alpha,\alpha^{'}),k \leftrightarrow c,(\gamma,\gamma^{'}),q$ we shall get:

$$\mathcal{V}_{abc}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) = \; tr([\lambda^{a},\lambda^{b}]\lambda^{c}) \; [\; F_{S\;\;\;\;\;A}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) + F_{A\;\;\;\;\;S}^{\alpha\alpha^{'}\beta\gamma\gamma^{'}}(k,p,q) \;] \tag{3.74}$$

which is in agreement with the expression we got in the tensor gauge field theory (2.25) and (2.26).

4 *Scattering Amplitude of Two Scalars and Two Tensors*

We shall get less trivial information about the structure of the VTT vertex of the open string theory if we shall compute the four particle scattering amplitude of two tachyon and two tensor bosons (see Fig.4). In the low energy limit these amplitudes will be dominated by the exchange of the massless vector boson (see Fig.5) and shall provide the information about the structure of cubic vertices $\mathcal{V}_{1-2-2}$ and $\mathcal{V}_{0-1-2}$. We are interested to calculate the following scattering amplitude on the disk (see Fig.4):

$$
\begin{gathered}
F^{\mu\nu,\lambda\rho}(k_1,k_2;k_3,k_4) = \\
= \int \prod_i \; d\mu(y_i) <: e^{ik_1X}(y_1) :: e^{ik_2X}(y_2) :: \dot{X}^\mu \dot{X}^\nu e^{ik_3X}(y_3) :: \dot{X}^\lambda \dot{X}^\rho e^{ik_4X}(y_4) :> \\
= \int \; d\mu(y_i) \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \\
\{ \; (F^\mu F^\nu)_{y_3} (F^\lambda F^\rho)_{y_4} + \\
+ (-2\alpha')[F^\mu_{y_3} F^\lambda_{y_4} \frac{\eta^{\nu\rho}}{(y_3-y_4)^2} + F^\mu_{y_3} F^\rho_{y_4} \frac{\eta^{\nu\lambda}}{(y_3-y_4)^2} + F^\nu_{y_3} F^\lambda_{y_4} \frac{\eta^{\mu\rho}}{(y_3-y_4)^2} + F^\nu_{y_3} F^\rho_{y_4} \frac{\eta^{\mu\lambda}}{(y_3-y_4)^2}] + \\
+ (-2\alpha')^2 [\frac{\eta^{\mu\lambda}}{(y_3-y_4)^2} \frac{\eta^{\nu\rho}}{(y_3-y_4)^2} + \frac{\eta^{\mu\rho}}{(y_3-y_4)^2} \frac{\eta^{\nu\lambda}}{(y_3-y_4)^2}]\}. \qquad (4.75)
\end{gathered}
$$

We shall fix the integration measure by the choice $y_4 = 0, y_2 = 1, y_1 \to \infty$. The mass-shell conditions are

$$
\alpha' k_1^2 = \alpha' k_2^2 = +1, \qquad \alpha' k_3^2 = \alpha' k_4^2 = -1
$$

and $k_1 + k_2 + k_3 + k_4 = 0$. Thus

$$
\begin{aligned}
\int \; d\mu(y_i) \prod_{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \; &= \int_{-\infty}^{+\infty} dy_3 \; y_1^2 \prod_{y_4=0,\; y_2=1,\; y_1\to\infty}^{i<j} |y_i - y_j|^{2\alpha' k_i k_j} \;\; \to \\
&= \int_{-\infty}^{+\infty} \; dy_3 \; |y_3|^{2\alpha' k_3 k_4} \; |1-y_3|^{2\alpha' k_2 k_3} \qquad (4.76)
\end{aligned}
$$

and

$$
\begin{aligned}
F^\mu_{y_3} &= -2i\alpha' \; (\frac{k_1^\mu}{y_3-y_1} + \frac{k_2^\mu}{y_3-y_2} + \frac{k_4^\mu}{y_3-y_4}) \to -2i\alpha' \; (\frac{k_2^\mu}{y_3-1} + \frac{k_4^\mu}{y_3}), \\
F^\mu_{y_4} &= -2i\alpha' \; (\frac{k_1^\mu}{y_4-y_1} + \frac{k_2^\mu}{y_4-y_2} + \frac{k_3^\mu}{y_4-y_3}) \to -2i\alpha' \; (-k_2^\mu - \frac{k_3^\mu}{y_3}). \qquad (4.77)
\end{aligned}
$$

Therefore we shall get

$$
\begin{gathered}
F^{\mu\nu,\lambda\rho}(k_1,k_2,k_3,k_4) = \int_{-\infty}^{+\infty} \; dy_3 \; |y_3|^{2\alpha' k_3 k_4} \; |1-y_3|^{2\alpha' k_2 k_3} \\
\{ \;\; (2\alpha')^4 (\frac{k_2^\mu}{y_3-1} + \frac{k_4^\mu}{y_3})(\frac{k_2^\nu}{y_3-1} + \frac{k_4^\nu}{y_3})(k_2^\lambda + \frac{k_3^\lambda}{y_3})(k_2^\rho + \frac{k_3^\rho}{y_3}) \\
- (2\alpha')^3 \; \frac{1}{y_3^2} \; [(\frac{k_2^\mu}{y_3-1} + \frac{k_4^\mu}{y_3})((k_2^\rho + \frac{k_3^\rho}{y_3})\eta^{\nu\lambda} + (k_2^\lambda + \frac{k_3^\lambda}{y_3})\eta^{\nu\rho})) \\
+ (\frac{k_2^\nu}{y_3-1} + \frac{k_4^\nu}{y_3})((k_2^\lambda + \frac{k_3^\lambda}{y_3})\eta^{\mu\rho} + (k_2^\rho + \frac{k_3^\rho}{y_3})\eta^{\mu\lambda})] \\
+ (2\alpha')^2 \; \frac{1}{y_3^4} \; [\eta^{\mu\lambda}\eta^{\nu\rho} + \eta^{\mu\rho}\eta^{\nu\lambda}] \;\; \} \qquad (4.78)
\end{gathered}
$$



Figure 4: Orderings of four open string vertex operators on the disk. The integral over y_3 in (4.76) splits into three ranges $[-\infty,0],[0,1],[1,+\infty]$. For these three ranges the vertex operators are ordered as in figures (d),(a) and (c) respectively. For the reversed cyclic permutation we shall get figures (f) (e) and (b). The coordinate y_3 increases in the direction of arrow.

The integration over y_3 can be divided into three regions $[-\infty,0],[0,1],[1,+\infty]$. These pieces can be depicted by three diagrams d) a) and c) on the Fig. 4. For the reversed cyclic permutation we shall get diagrams f) e) and b) on the Fig. 4. Introducing Mandelshtam variables

$$s=-(k_1+k_2)^2, \quad t=-(k_2+k_3)^2, \quad u=-(k_2+k_4)^2$$

we can represent the contribution of the (s,t) diagrams a) and e) in the form

$$\begin{aligned} &-(2\alpha')^3\ (tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_3}\lambda^{a_4})+tr(\lambda^{a_4}\lambda^{a_3}\lambda^{a_2}\lambda^{a_1}))\\ &[B(-\alpha' s,-\alpha' t+1)K^{\mu\rho\nu\lambda}(k_4,k_2)+B(-\alpha' s-1,-\alpha' t+1)K^{\mu\rho\nu\lambda}(k_4,k_3)\\ &-B(-\alpha' s+1,-\alpha' t)K^{\mu\rho\nu\lambda}(k_2,k_2)-B(-\alpha' s,-\alpha' t)K^{\mu\rho\nu\lambda}(k_2,k_3)], \end{aligned} \tag{4.79}$$

the contribution of the (s,u) diagrams f) and c) in the form

$$\begin{aligned} &-(2\alpha')^3\ (tr(\lambda^{a_1}\lambda^{a_2}\lambda^{a_4}\lambda^{a_3})+tr(\lambda^{a_1}\lambda^{a_3}\lambda^{a_4}\lambda^{a_2}))\\ &[-B(-\alpha' s,-\alpha' u)K^{\mu\rho\nu\lambda}(k_4,k_2)+B(-\alpha' s-1,-\alpha' u+1)K^{\mu\rho\nu\lambda}(k_4,k_3)\\ &-B(-\alpha' s+1,-\alpha' u)K^{\mu\rho\nu\lambda}(k_2,k_2)+B(-\alpha' s,-\alpha' u+1)K^{\mu\rho\nu\lambda}(k_2,k_3)] \end{aligned} \tag{4.80}$$

and the the contribution of the (u,t) diagrams b) and d) in the form

$$\begin{aligned} &-(2\alpha')^3\ (tr(\lambda^{a_1}\lambda^{a_3}\lambda^{a_2}\lambda^{a_4})+tr(\lambda^{a_1}\lambda^{a_4}\lambda^{a_2}\lambda^{a_3}))\\ &[B(-\alpha' u,-\alpha' t+1)K^{\mu\rho\nu\lambda}(k_4,k_2)+B(-\alpha' u+1,-\alpha' t+1)K^{\mu\rho\nu\lambda}(k_4,k_3)\\ &+B(-\alpha' u,-\alpha' t)K^{\mu\rho\nu\lambda}(k_2,k_2)+B(-\alpha' u+1,-\alpha' t)K^{\mu\rho\nu\lambda}(k_2,k_3)], \end{aligned} \tag{4.81}$$



Figure 5: Feynman diagrams contributing to the low-energy limit of the open string scattering amplitude. The poles correspond to the exchange by a massless vector boson in s,t and u channels. The scalars are depicted as 0, vectors as 1 and tensors as 2. The quantities of main interest here are the dimensionless vertices $\mathcal{V}_{1-2-2}$ between a vector and two tensors in s-channel and $\mathcal{V}_{0-1-2}$ between a scalar, vector and a tensor in t,u channels. The last graph represents the contact vertex $\mathcal{V}_{0-0-2-2}$.

where

$$K^{\mu\rho\nu\lambda}(k,p) = k^{\mu}(p^{\rho}\eta^{\nu\lambda} + p^{\lambda}\eta^{\nu\rho}) + k^{\nu}(p^{\lambda}\eta^{\mu\rho} + p^{\rho}\eta^{\mu\lambda}). \tag{4.82}$$

Considering the limit $\alpha' s, \alpha' t, \alpha' u \to 0$ of the Euler functions we shall get for the s , t and u channel contributions:

$$\begin{aligned} -(2\alpha')^3 \{ \ &+ \frac{1}{\alpha' s} tr[\lambda^{a_1},\lambda^{a_2}][\lambda^{a_3}\lambda^{a_4}](K(k_2,k_3) - K(k_4,k_2))^{\mu\rho\nu\lambda} \\ &- \frac{1}{\alpha' t} tr[\lambda^{a_1},\lambda^{a_4}][\lambda^{a_2}\lambda^{a_3}](K(k_2,k_2) + K(k_2,k_3))^{\mu\rho\nu\lambda} \\ &- \frac{1}{\alpha' u} tr[\lambda^{a_1},\lambda^{a_3}][\lambda^{a_2}\lambda^{a_4}](K(k_2,k_2) - K(k_4,k_2))^{\mu\rho\nu\lambda}\}. \end{aligned} \tag{4.83}$$

which are shown on the Fig.5.

The last term in the matrix element (4.78) has no momentum dependence and represents the contact term. Evaluating the integration over y_3 in same way as we did above one can get:

$$\begin{aligned} (2\alpha')^2 \{ \ &+ (\ tr\lambda^{a_1}\lambda^{a_2}\lambda^{a_3}\lambda^{a_4} + tr\lambda^{a_4}\lambda^{a_3}\lambda^{a_2}\lambda^{a_1}\ )B(-\alpha' s - 1, -\alpha' t + 1) \\ &+ (\ tr\lambda^{a_1}\lambda^{a_2}\lambda^{a_4}\lambda^{a_3} + tr\lambda^{a_1}\lambda^{a_3}\lambda^{a_4}\lambda^{a_2}\ )B(-\alpha' s - 1, -\alpha' u + 1) \\ &+ (\ tr\lambda^{a_1}\lambda^{a_3}\lambda^{a_2}\lambda^{a_4}) + tr\lambda^{a_1}\lambda^{a_4}\lambda^{a_2}\lambda^{a_3}\ )B(-\alpha' t + 1, -\alpha' u + 1)\} \\ &(\eta^{\mu\lambda}\eta^{\nu\rho} + \eta^{\mu\rho}\eta^{\nu\lambda}). \end{aligned} \tag{4.84}$$

Again considering the limit $\alpha' s, \alpha' t, \alpha' u \to 0$ of the Euler functions we shall get for the contact term

$$(2\alpha')^2 \ \{\ tr[\lambda^{a_1},\lambda^{a_4}][\lambda^{a_2},\lambda^{a_3}] + \frac{t}{s}\ tr[\lambda^{a_1},\lambda^{a_2}][\lambda^{a_3},\lambda^{a_4}]\ \}(\eta^{\mu\lambda}\eta^{\nu\rho} + \eta^{\mu\rho}\eta^{\nu\lambda}). \tag{4.85}$$

This is the quartic vertex of $\mathcal{V}_{0-0-2-2}$ and should be compared with correspond vertex in non-Abelian tensor gauge field theory [27, 28, 29].

5 *Conclusion*

Our intention in this article was to compare the structure of the tree level scattering amplitudes in non-Abelian tensor gauge field theory and in open string theory with Chan-Paton charges. We limit ourselves considering only lower rank tensor fields in both theories. We identify the symmetric T_S and antisymmetric T_A components of the second rank tensor gauge field $A^a_{\alpha\acute{\alpha}}$ with the string excitations T_S and T_A on the third and forth levels. In the process of this identification we have been selecting only those parts of the tree level scattering amplitudes in the open string theory which have *dimensionless coupling constants in four dimensions*. It should be mentioned that not all amplitudes have these dimensionless parts. In particular, the scattering amplitude of three tensor bosons T does not have dimensionless parts. It seems that *dimensionless subclass of tree level scattering amplitudes may provide information about the structure of the open string theory at the deepest level.*

In this respect one could mention that the cubic and quartic vertices of the massless Yang-Mills theory and a massive theory with spontaneous symmetry breaking are identical. Therefore if we want to extract a genuine massless proto-theory from the open string theory, which in accordance to David Gross is in a broken phase [4], it seems very natural to consider a subclass of amplitudes with dimensionless coupling constant and compare them with the amplitudes in non-Abelian tensor gauge field theory.

It is also interesting to mention that the ratio of the masses of the tensor gauge bosons T_S and T_A in non-Abelian tensor gauge theory with spontaneous symmetry breaking is [54]

$$\frac{m_A^2}{m_S^2} = 3, \tag{5.86}$$

while in the open string theory it is (see Fig.1)

$$\frac{m_A^2}{m_S^2} = 4. \tag{5.87}$$

In both theories the antisymmetric tensor is more heavy.

In conclusion I would like to thank Costas Bachas for stimulating discussions and his suggestion of comparing scattering amplitudes in both theories. I also would like to thank Sebastian Guttenberg and Spiros Konitopoulos for many discussions and CERN Theory Division for hospitality, where part of this work was completed. This work was partially supported by ENRAGE (European Network on Random Geometry), Marie Curie Research Training Network, contract MRTN-CT-2004- 005616.

References

[1] A. Neveu and J. Scherk, *Connection between Yang-Mills fields and dual models,* Nucl. Phys. B **36** (1972) 155.

[2] M. B. Green, J. H. Schwarz and E. Witten, *Superstring theory. Vol. 1: Introduction, Cambridge, UK: Univ. Pr. (1987) 469 P. (Cambridge Monographs On Mathematical Physics)*

[3] J. Polchinski, *String theory. Vol. 1: An introduction to the bosonic string, Cambridge, UK: Univ. Pr. (1998) 402 p*

[4] D. J. Gross, *High-Energy Symmetries Of String Theory*, Phys. Rev. Lett. **60** (1988) 1229.

[5] D. J. Gross and P. F. Mende, *The High-Energy Behavior Of String Scattering Amplitudes*, Phys. Lett. B **197** (1987) 129.

[6] D. J. Gross and P. F. Mende, *String Theory Beyond The Planck Scale*, Nucl. Phys. B **303** (1988) 407.

[7] E. Witten, *The Search For Higher Symmetry In String Theory*, Philos. Trans. R. Soc. London A320 (1989) 349-357;
E. Witten, *Space-Time And Topological Orbifolds*, Phys. Rev. Lett. **61** (1988) 670

[8] P. F. Mende, *String theory at short distance and the principle of equivalence*, arXiv:hep-th/9210001.

[9] P. F. Mende and H. Ooguri, *Borel Summation Of String Theory For Planck Scale Scattering*, Nucl. Phys. B **339** (1990) 641.

[10] P. F. Mende, *High-energy string collisions in a compact space*, Phys. Lett. B **326** (1994) 216

[11] G. W. Moore, *Symmetries of the bosonic string S matrix*, arXiv:hep-th/9310026.

[12] G. K. Savvidy, "Tensionless strings: Physical Fock space and higher spin fields," Int. J. Mod. Phys. A **19**, (2004) 3171-3194.

[13] S. B. Giddings, D. J. Gross and A. Maharana, *Gravitational effects in ultrahigh-energy string scattering,* arXiv:hep-th/0705.1816

[14] G. Savvidy, *Tensionless strings, correspondence with SO(D,D) sigma model*, Phys. Lett. B **615** (2005) 285.

[15] D. Amati, M. Ciafaloni and G. Veneziano, *Can Space-Time Be Probed Below The String Size?* Phys. Lett. B **216** (1989) 41.

[16] U. Lindstrom, B. Sundborg and G. Theodoridis, *The Zero Tension Limit Of The Superstring,* Phys. Lett. B **253** (1991) 319.

[17] H. J. De Vega and A. Nicolaidis, *Strings in strong gravitational fields,* Phys. Lett. B **295** (1992) 214.

[18] I. Bakas and C. Sourdis, *On the tensionless limit of gauged WZW models,* arXiv:hep-th/0403165.

[19] E. Witten, *Noncommutative Geometry And String Field Theory,* Nucl. Phys. B **268** (1986) 253.

[20] C. B. Thorn, *Comments On Covariant Formulations Of String Theories,* Phys. Lett. **159B** (1985) 107; *String Field Theory,* Phys. Rep. **174C** (1989) 1

[21] W. Siegel and B. Zwiebach, *Gauge String Fields,* Nucl. Phys. B **263** (1986) 105.

[22] I. Y. Arefeva, P. B. Medvedev and A. P. Zubarev, *New Representation For String Field Solves The Consistence Problem For Open Superstring Field,* Nucl. Phys. B **341** (1990) 464.

[23] W. Siegel, *Introduction to string field theory,* arXiv:hep-th/0107094.

[24] W. Taylor and B. Zwiebach, *D-branes, tachyons, and string field theory,* arXiv:hep-th/0311017 (see formulas (201) and (202) of section 6.5).

[25] W. Taylor, *String field theory,* arXiv:hep-th/0605202.

[26] C.N.Yang and R.L.Mills. "Conservation of Isotopic Spin and Isotopic Gauge Invariance". Phys. Rev. **96** (1954) 191

[27] G. Savvidy, *Non-Abelian tensor gauge fields: Generalization of Yang-Mills theory,* Phys. Lett. B **625** (2005) 341

[28] G. Savvidy, *Non-abelian tensor gauge fields. I,* Int. J. Mod. Phys. A **21** (2006) 4931;

[29] G. Savvidy, *Non-abelian tensor gauge fields. II,* Int. J. Mod. Phys. A **21** (2006) 4959;

[30] G. Savvidy and T. Tsukioka, *Gauge invariant Lagrangian for non-Abelian tensor gauge fields of fourth rank,* Prog. Theor. Phys. **117** (2007) 4; arXiv:hep-th/0512344.

[31] J. Barrett and G. Savvidy *A Dual Lagrangian for Non-Abelian Tensor Gauge Fields*, Phys. Lett. B **652** (2007) 141-145

[32] S. Guttenberg and G. Savvidy, *Duality transformation of non-Abelian tensor gauge fields,* Mod. Phys. Lett. A **23** (2008) 999

[33] S. Konitopoulos, R. Fazio and G. Savvidy, *Tensor gauge boson production in high energy collisions,* arXiv:0803.0075 [hep-th].

[34] S. Konitopoulos and G. Savvidy, *Production of Spin-Two Gauge Bosons,* arXiv:0804.0847 [hep-th].

[35] J. E. Paton and H. M. Chan, *Generalized veneziano model with isospin,* Nucl. Phys. B **10** (1969) 516.

[36] H. Kawai, D. C. Lewellen and S. H. H. Tye, *A Relation Between Tree Amplitudes Of Closed And Open Strings,* Nucl. Phys. B **269** (1986) 1.

[37] F. A. Berends, R. Kleiss, P. De Causmaecker, R. Gastmans and T. T. Wu, *Single Bremsstrahlung Processes In Gauge Theories,* Phys. Lett. B **103** (1981) 124.

[38] R. Kleiss and W. J. Stirling, *Spinor Techniques For Calculating P Anti-P → W+- / Z0 + Jets,* Nucl. Phys. B **262** (1985) 235.

[39] Z. Xu, D. H. Zhang and L. Chang, *Helicity Amplitudes for Multiple Bremsstrahlung in Massless Nonabelian Gauge Theories,* Nucl. Phys. B **291** (1987) 392.

[40] J. F. Gunion and Z. Kunszt, *Improved Analytic Techniques For Tree Graph Calculations And The G G Q Anti-Q Lepton Anti-Lepton Subprocess,* Phys. Lett. B **161** (1985) 333.

[41] L. J. Dixon, *Calculating scattering amplitudes efficiently,* arXiv:hep-ph/9601359.

[42] S. J. Parke and T. R. Taylor, *An Amplitude for n Gluon Scattering,* Phys. Rev. Lett. **56** (1986) 2459.

[43] F. A. Berends and W. T. Giele, *Recursive Calculations for Processes with n Gluons,* Nucl. Phys. B **306** (1988) 759.

[44] E. Witten, *Perturbative gauge theory as a string theory in twistor space,* Commun. Math. Phys. **252** (2004) 189 [arXiv:hep-th/0312171].

[45] R. Britto, F. Cachazo and B. Feng, *New recursion relations for tree amplitudes of gluons,* Nucl. Phys. B **715** (2005) 499 [arXiv:hep-th/0412308].

[46] R. Britto, F. Cachazo, B. Feng and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory,* Phys. Rev. Lett. **94** (2005) 181602 [arXiv:hep-th/0501052].

[47] P. Benincasa and F. Cachazo, *Consistency Conditions on the S-Matrix of Massless Particles,* arXiv:0705.4305 [hep-th].

[48] R. R. Metsaev, *Cubic interaction vertices for fermionic and bosonic arbitrary spin fields,* arXiv:0712.3526 [hep-th].

[49] T. Curtright, *Generalized Gauge Fields,* Phys. Lett. B **165** (1985) 304.

[50] A. K. Bengtsson, I. Bengtsson and L. Brink, *Cubic Interaction Terms For Arbitrary Spin,* Nucl. Phys. B **227** (1983) 31.

[51] A. K. Bengtsson, I. Bengtsson and L. Brink, *Cubic Interaction Terms For Arbitrarily Extended Supermultiplets,* Nucl. Phys. B **227** (1983) 41.

[52] A. K. H. Bengtsson, I. Bengtsson and N. Linden, *Interacting Higher Spin Gauge Fields on the Light Front,* Class. Quant. Grav. **4** (1987) 1333.

[53] R. R. Metsaev, *Cubic interaction vertices of massive and massless higher spin fields,* Nucl. Phys. B **759** (2006) 147

[54] G. Savvidy, *Interaction of non-Abelian tensor gauge fields,* arXiv:0804.2003 [hep-th].